

Question 1:

(a)[3 points] Evaluate $f'(0)$ where $f(x) = x \arccos(x) - \sqrt{1-x^2}$.

$$f'(x) = \arccos(x) + x \left(\frac{-1}{\sqrt{1-x^2}} \right) - \frac{1}{2} (1-x^2)^{-\frac{1}{2}} (-2x)$$

$$f'(0) = \arccos(0)$$

$$= \boxed{\frac{\pi}{2}}$$

(b)[3 points] Evaluate $g'(0)$ where $g(x) = \cosh(x) \sinh(x^2)$.

$$g'(x) = \sinh(x) \sinh(x^2) + \cosh(x) \cosh(x^2) (2x)$$

$$\therefore g'(0) = \boxed{0}$$

(c)[4 points] Evaluate

$$\lim_{x \rightarrow 0^+} \sin(x) \ln(x) \sim "0 \cdot (-\infty)"$$

$$= \lim_{x \rightarrow 0^+} \frac{\ln x}{\left(\frac{1}{\sin(x)}\right)} \sim \frac{-\infty}{\infty}$$

$$\stackrel{H}{=} \lim_{x \rightarrow 0^+} \frac{\frac{1}{x}}{\left(\frac{-1}{\sin^2 x}\right) \cos x}$$

$$= \lim_{x \rightarrow 0^+} -\frac{\sin^2 x}{x \cos x} \sim \frac{0}{0}$$

$$\stackrel{H}{=} \lim_{x \rightarrow 0^+} \frac{-2 \sin x \cos x}{\cos x - x \sin x}$$

$$= \boxed{0}$$

Question 2:

(a) [4 points] Evaluate

$$\begin{aligned} \lim_{t \rightarrow 0} \left(\frac{1}{t} - \frac{1}{te^t} \right) &\sim \text{"} \pm (\infty - \infty) \text{"} \\ &= \lim_{t \rightarrow 0} \frac{e^t - 1}{te^t} \sim \frac{0}{0} \\ &\stackrel{H}{=} \lim_{t \rightarrow 0} \frac{e^t}{e^t + te^t} \\ &= \lim_{t \rightarrow 0} \frac{1}{1+t} \\ &= \boxed{1} \end{aligned}$$

(b) [3 points] An animal gains mass at a rate of $W(t) = \frac{100t}{(t^2+1)^3}$ kg/yr, where t is time in years. What is the total mass gain during the first two years of life?

Let $w(t)$ = mass at time t years, so $w'(t) = W(t)$.

$$\begin{aligned} w(2) - w(0) &= \int_0^2 \frac{100t}{(t^2+1)^3} dt \quad \begin{cases} u = t^2+1, du = 2t dt \\ t=0 \Rightarrow u=1 \\ t=2 \Rightarrow u=5 \end{cases} \\ &= 50 \int_1^5 u^{-3} du \\ &= \frac{50}{-2} [u^{-2}]_1^5 \\ &= -25 \left[\frac{1}{25} - 1 \right] = \boxed{24 \text{ kg}} \end{aligned}$$

(c) [3 points] The average value of $f(x) = qx^2$ over the interval $[-q, q]$ is 9. Determine the value of the constant q .

$$\begin{aligned} 9 &= \frac{1}{q - (-q)} \int_{-q}^q qx^2 dx \\ &= \frac{1}{2q} \cdot q \cdot \left[\frac{x^3}{3} \right]_{-q}^q \\ \therefore 18 &= \frac{q^3}{3} - \left(-\frac{q^3}{3} \right) \\ 18 &= \frac{2}{3} q^3 \\ \therefore q &= \left[\frac{(3)(18)}{2} \right]^{\frac{1}{3}} = \boxed{3} \end{aligned}$$

Question 3:

(a)[3 points] Define the function

$$F(x) = \int_{-1}^{x^3} \frac{\cos(t^2)}{e^t} dt$$

Evaluate and simplify $F(-1) - F'(0)$.

$$F(-1) = \int_{-1}^{(-1)^3} \frac{\cos(t^2)}{e^t} dt = \int_{-1}^{-1} \frac{\cos(t^2)}{e^t} dt = 0$$

$$F'(x) = \frac{\cos((x^3)^2)}{e^{x^3}} \cdot 3x^2$$

$$\therefore F'(0) = 0$$

$$\therefore F(-1) - F'(0) = \boxed{0}$$

(b)[3 points] Evaluate:

$$I = \int \frac{1}{x \ln x} dx$$

$$\text{let } u = \ln x, \quad du = \frac{1}{x} dx$$

$$\therefore I = \int \frac{1}{u} du$$

$$= \ln|u| + C$$

$$= \boxed{\ln|\ln x| + C}$$

(c)[4 points] Evaluate:

$$\int_0^1 (\sqrt{x} + 1)^2 dx$$

$$= \int_0^1 x^{\frac{1}{2}} + 2x^{\frac{1}{4}} + 1 dx$$

$$= \left[\frac{2}{3} x^{\frac{3}{2}} + 2\left(\frac{4}{5}\right) x^{\frac{5}{4}} + x \right]_0^1$$

$$= \frac{2}{3} + \frac{8}{5} + 1$$

$$= \frac{10 + 24 + 15}{15}$$

$$= \boxed{\frac{49}{15}}$$

Question 4:

(a)[3 points] Evaluate:

$$I = \int \frac{x+2}{(x-1)^{3/2}} dx$$

$$\text{let } u = x-1 \\ du = dx$$

$$\therefore I = \int \frac{u+3}{u^{3/2}} du$$

$$= \int u^{-1/2} + 3u^{-3/2} du$$

$$= 2u^{1/2} - 3(2)u^{-1/2} + C$$

$$= \boxed{2(x-1)^{1/2} - 6(x-1)^{-1/2} + C}$$

(b)[4 points] Evaluate the following limit of Riemann Sums by interpreting it as $\int_a^b f(x) dx$ for some a , b and $f(x)$:


$$I = \lim_{n \rightarrow \infty} \frac{1}{n} \left[\left(\frac{1}{n}\right)^3 + \left(\frac{2}{n}\right)^3 + \dots + \left(\frac{n}{n}\right)^3 \right]$$

$$= \lim_{n \rightarrow \infty} \sum_{i=1}^n \left(\frac{i}{n}\right)^3 \left(\frac{1}{n}\right)$$

$$\therefore x_i = \frac{i}{n}, \quad \Delta x = \frac{1}{n}, \quad f(x_i) = (x_i)^3$$

$$\therefore I = \int_0^1 x^3 dx$$

$$= \left[\frac{x^4}{4} \right]_0^1 = \boxed{\frac{1}{4}}$$

(c)[3 points] For the function $f(x)$ whose graph is shown below, $\int_{-4}^5 f(x) dx = \frac{-5}{6}$. Determine the area of the shaded region A .

$$\therefore \int_{-4}^{-3} f(x) dx + \int_{-3}^3 f(x) dx + \int_3^5 f(x) dx = \frac{-5}{6}$$

$$\frac{1}{2} + 0 - A = \frac{-5}{6}$$

$$\therefore A = \frac{1}{2} + \frac{5}{6} = \frac{3+5}{6} = \boxed{\frac{4}{3}}$$

Question 5 [8 points]: Evaluate:

$$I = \int x^3 (\ln x)^2 dx$$

$$u = (\ln x)^2 ; dv = x^3$$

$$du = 2 \ln x \cdot \frac{1}{x} ; v = \frac{x^4}{4}$$

$$\therefore I = (\ln x)^2 \frac{x^4}{4} - \int \frac{x^4}{4} \cdot 2 \ln x \cdot \frac{1}{x} dx$$

$$= \frac{(\ln x)^2 x^4}{4} - \frac{1}{2} \int x^3 \ln x dx$$

$$u = \ln x ; dv = x^3$$

$$du = \frac{1}{x} dx ; v = \frac{x^4}{4}$$

$$= \frac{(\ln x)^2 x^4}{4} - \frac{1}{2} \left[(\ln x) \left(\frac{x^4}{4} \right) - \int \frac{x^4}{4} \frac{1}{x} dx \right]$$

$$= \frac{(\ln x)^2 x^4}{4} - \frac{1}{8} (\ln x) x^4 + \frac{1}{8} \int x^3 dx$$

$$= \frac{(\ln x)^2 x^4}{4} - \frac{1}{8} (\ln x) x^4 + \frac{1}{32} x^4 + C$$

Question 6 [8 points]: Evaluate:

$$I = \int \frac{5x-8}{x^2+x-12} dx$$

$$\begin{aligned} \frac{5x-8}{x^2+x-12} &= \frac{5x-8}{(x+4)(x-3)} = \frac{A}{x+4} + \frac{B}{x-3} \\ &= \frac{A(x-3) + B(x+4)}{(x+4)(x-3)} \\ &= \frac{(A+B)x - 3A + 4B}{(x+4)(x-3)} \end{aligned}$$

$$\begin{aligned} \therefore A+B &= 5 \Rightarrow B = 5-A \\ -3A+4B &= -8 \Rightarrow -3A+4(5-A) = -8 \\ &\quad -7A+20 = -8 \\ &\quad -7A = -28 \\ &\quad \therefore A = 4 \\ &\quad \therefore B = 5-4 = 1 \end{aligned}$$

$$\therefore I = \int \frac{4}{x+4} + \frac{1}{x-3} dx$$

$$= \boxed{4 \ln|x+4| + \ln|x-3| + C}$$

Question 7 [8 points]: Evaluate:

$$I = \int \sqrt{16-x^2} dx$$

$$x = 4 \sin \theta$$

$$dx = 4 \cos \theta d\theta$$

$$\therefore I = \int \sqrt{16-16\sin^2\theta} 4\cos\theta d\theta$$


$$= \int 4\sqrt{1-\sin^2\theta} 4\cos\theta d\theta$$

$$= 16 \int \cos^2\theta d\theta$$

$$= 16 \int \frac{1 + \cos(2\theta)}{2} d\theta$$

$$= 16 \left[\frac{\theta}{2} + \frac{\sin(2\theta)}{4} \right] + C$$

$$= 8\theta + 8\sin\theta\cos\theta + C$$



$$\therefore \sin\theta = \frac{x}{4}$$

$$\cos\theta = \frac{\sqrt{16-x^2}}{4}$$


$$\theta = \arcsin\left(\frac{x}{4}\right)$$

$$\therefore I = 8\arcsin\left(\frac{x}{4}\right) + 8\left(\frac{x}{4}\right)\frac{\sqrt{16-x^2}}{4} + C$$

$$= 8\arcsin\left(\frac{x}{4}\right) + \frac{1}{2} \cdot x \cdot \sqrt{16-x^2} + C$$

Question 8:

- (a)[4 points] Consider the integral $\int_0^4 4(1+x^2) dx$. Let M_2 be the midpoint rule approximation of the integral using two subintervals, and T_2 the trapezoid rule approximation using two subintervals. Determine $T_2 - M_2$.



$$T_2 = \left(\frac{4+20}{2}\right)(2) + \left(\frac{20+68}{2}\right)(2) = 112$$

$$M_2 = (8)(2) + (40)(2) = 96$$

$$\therefore T_2 - M_2 = 112 - 96 = \boxed{16}$$

- (b)[4 points] Determine the error in T_2 , the trapezoid rule approximation used in part (a). Recall, the error in using the trapezoid rule to approximate $\int_a^b f(x) dx$ is at most $\frac{K(b-a)^3}{12n^2}$, where $|f''(x)| \leq K$ on $[a, b]$.

$$f(x) = 4 + 4x^2$$

$$f'(x) = 8x$$

$$f''(x) = 8 = K$$

$$\therefore |E_{T_2}| \leq \frac{8(4-0)^3}{12 \cdot 2^2} = \frac{8 \cdot 4 \cdot 4 \cdot 4}{3 \cdot 4 \cdot 4} = \boxed{\frac{32}{3}}$$

Question 9:

(a)[4 points] Evaluate the improper integral

$$\begin{aligned}
 & \int_0^{\infty} \frac{x^3}{e^{(x^4)}} dx \\
 &= \lim_{b \rightarrow \infty} \int_0^b x^3 e^{-x^4} dx \\
 &= \lim_{b \rightarrow \infty} \left[\frac{e^{-x^4}}{-4} \right]_0^b \\
 &= \lim_{b \rightarrow \infty} \left[\frac{e^{-b^4}}{-4} + \frac{1}{4} \right] \\
 &= \boxed{\frac{1}{4}}
 \end{aligned}$$

(b)[4 points] Evaluate the improper integral:

$$\begin{aligned}
 & \int_2^3 \frac{1}{\sqrt{3-x}} dx \\
 &= \lim_{b \rightarrow 3^-} \int_2^b [3-x]^{-\frac{1}{2}} dx \\
 &= \lim_{b \rightarrow 3^-} \left[-2[3-x]^{\frac{1}{2}} \right]_2^b \\
 &= \lim_{b \rightarrow 3^-} \left[-2(3-b)^{\frac{1}{2}} + 2(3-2)^{\frac{1}{2}} \right] \\
 &= \boxed{2}
 \end{aligned}$$

(c)[2 points] Use the Comparison Theorem to show that the following integral diverges:


$$\int_1^{\infty} \frac{x^3 + \sin^2 x}{x^4} dx$$

$$\frac{x^3 + \sin^2 x}{x^4} \geq \frac{x^3}{x^4} = \frac{1}{x}$$

Since $\int_1^{\infty} \frac{1}{x} dx$ diverges, by the Comparison Theorem, so must $\int_1^{\infty} \frac{x^3 + \sin^2 x}{x^4} dx$.

Question 10:

- (a) [5 points] Determine the area of the region bounded between the curves $y = 4x - x^2$ and $y = x^2$.




$$A = \int_0^2 (4x - x^2 - x^2) dx$$

$$= \left[\frac{4x^2}{2} - \frac{2x^3}{3} \right]_0^2$$

$$= 8 - \frac{16}{3}$$

$$= \frac{24-16}{3} = \boxed{\frac{8}{3}}$$

- (b) [5 points] The bounded region in part (a) is rotated about the x -axis. Determine the volume of the resulting solid. (Disks/washers would be best here.)



$$V = \int_0^2 \pi \left[(4x - x^2)^2 - (x^2)^2 \right] dx$$


$$= \pi \int_0^2 \left[16x^2 - 8x^3 + \cancel{x^4} - \cancel{x^4} \right] dx$$

$$= \pi \left[\frac{16x^3}{3} - \frac{8x^4}{4} \right]_0^2$$

$$= \pi \left[\frac{128}{3} - 32 \right] = \pi \left[\frac{128-96}{3} \right] = \boxed{\pi \cdot \frac{32}{3}}$$


Question 11:

- (a) [4 points] The curve $y = \cos(x)$, $-\pi/2 \leq x \leq \pi/2$, is rotated about the vertical line $x = \pi$. Set up but do not evaluate the integral representing the volume of the resulting solid. (Cylinders would be best here.)



$$V = \int_{-\pi/2}^{\pi/2} 2\pi (\pi - x) \cos(x) dx$$

- (b) [4 points] 20 m rope hangs over the side of a building and a 10 kg bucket is tied to the end of the rope. A person at the top of the building pulls the rope and bucket up onto the roof of the building. How much work is done if the rope has a total mass of 2 kg? Recall that the acceleration due to gravity is $g = 9.8 \text{ m/s}^2$.



$$W_{\text{bucket}} = (10 \text{ kg}) \left(9.8 \frac{\text{m}}{\text{s}^2} \right) (20 \text{ m})$$

$$= 1960 \text{ Nm}$$

$$W_{\text{rope}} = \int_{y=0}^{20} \left(\frac{2 \text{ kg}}{20 \text{ m}} \right) (y \text{ m}) \left(9.8 \frac{\text{m}}{\text{s}^2} \right) (dy \text{ m})$$

$$= \frac{9.8}{10} \int_0^{20} y dy$$

$$= \frac{9.8}{10} \left[\frac{y^2}{2} \right]_0^{20}$$


$$= \frac{9.8}{10} \cdot \frac{20}{2} \cdot 20$$

$$= 196 \text{ Nm}$$

$$\therefore \text{Total work is } W_{\text{bucket}} + W_{\text{rope}} = 1960 + 196 = \boxed{2156 \text{ Nm}}$$

Question 12:

- (a) [5 points] The base (flat bottom surface) of a solid is the region between the curve $y = \sqrt{1-x^2}$ and the x -axis. Note that $y = \sqrt{1-x^2}$ is the upper half of the circle of radius 1 and centre $(0,0)$. Cross-sections perpendicular to the x -axis are isosceles triangles of equal height and base. Determine the volume of the solid.



$$\begin{aligned} \therefore A(x) &= \frac{1}{2} \sqrt{1-x^2} \cdot \sqrt{1-x^2} \\ &= \frac{1-x^2}{2} \end{aligned}$$

$$\begin{aligned} \therefore V &= \int_{-1}^1 \frac{1-x^2}{2} dx \\ &= \frac{1}{2} \left[x - \frac{x^3}{3} \right]_{-1}^1 \\ &= \frac{1}{2} \left[\left(1 - \frac{1}{3}\right) - \left(-1 + \frac{1}{3}\right) \right] \\ &= \boxed{\frac{2}{3}} \end{aligned}$$

- (b) [5 points] Solve the following differential equation:

$$\frac{dy}{dx} = \frac{1+y^2}{\sqrt{x+1}}, \quad y(0) = 1$$

You may leave your solution in implicit form (it is not necessary to isolate the y variable in your final answer.)

$$\int \frac{1}{1+y^2} dy = \int (x+1)^{-\frac{1}{2}} dx$$

$$\arctan(y) = 2(x+1)^{\frac{1}{2}} + C$$

$$y(0) = 1 :$$

$$\arctan(1) = 2 + C$$

$$\therefore C = \frac{\pi}{4} - 2$$

$$\therefore \boxed{\arctan(y) = 2(x+1)^{\frac{1}{2}} + \frac{\pi}{4} - 2}$$

Question 13:

(a) [3 points] Determine the first three non-zero terms of the Maclaurin series for the function $f(x) = x^3 e^{x^2}$.

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots$$

$$e^{x^2} = 1 + x^2 + \frac{x^4}{2!} + \frac{x^6}{3!} + \dots$$

$$x^3 e^{x^2} = x^3 + x^5 + \frac{x^7}{2!} + \dots$$

\therefore first three non-zero terms are $x^3 + x^5 + \frac{x^7}{2!}$

(b) [4 points] Use a Maclaurin series (not L'Hospital's Rule) to evaluate the limit

$$\begin{aligned} & \lim_{x \rightarrow 0} \frac{1 - \cos(x^2)}{x^3(e^x - 1)} \\ &= \lim_{x \rightarrow 0} \frac{1 - \left[1 - \frac{x^4}{2!} + \frac{x^8}{3!} - \dots \right]}{x^3 \left[1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots - 1 \right]} \\ &= \lim_{x \rightarrow 0} \frac{\frac{x^4}{2!} - \frac{x^8}{3!} + \dots}{x^4 + \frac{x^5}{2!} + \frac{x^6}{3!}} \\ &= \lim_{x \rightarrow 0} \frac{\frac{1}{2!} - \frac{x^4}{3!} + \dots}{1 + \frac{x}{2!} + \frac{x^2}{3!} + \dots} \\ &= \boxed{\frac{1}{2}} \end{aligned}$$

(c) [3 points] Determine the Maclaurin polynomial of degree three for $f(x) = e^x \sin x$. You may use the definition or any other valid method to obtain your answer.

$$\begin{aligned} f(x) &= \left(1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots \right) \left(x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots \right) \\ &= x + x^2 - \frac{x^3}{3!} + \frac{x^3}{2!} + \left(\text{terms of degree greater than 3} \right) \\ &= x + x^2 + \frac{3x^3 - x^3}{6} + \left(\text{terms of degree greater than 3} \right) \\ &\therefore \boxed{T_3(x) = x + x^2 + \frac{1}{3} x^3} \end{aligned}$$