

Math 370 - Introductory Complex Variables

G.Pugh

Nov 13 2014

Recap of Last Day

- ▶ **Theorem:** If f is analytic inside and on the simple closed positively oriented contour Γ and z is inside Γ then

$$f^{(n)}(z) = \frac{n!}{2\pi i} \int_{\Gamma} \frac{f(\zeta)}{(\zeta - z)^{n+1}} d\zeta$$

- ▶ Consequently: If f is analytic inside and on Γ , then $f'(z)$, $f''(z)$, $f'''(z)$, \dots all exist for z inside Γ .

Liouville's Theorem

Theorem: A bounded entire function is a constant.

Proof: Suppose that f is entire and that $|f(z)| \leq M$ for every $z \in \mathbb{C}$. Let $z \in \mathbb{C}$ and let C_R be any circle of radius R , centre z . Then

$$f'(z) = \frac{1}{2\pi i} \int_{C_R} \frac{f(\zeta)}{(\zeta - z)^2} d\zeta$$

On C_R , $\left| \frac{f(\zeta)}{(\zeta - z)^2} \right| \leq \frac{M}{R^2}$, and the length of C_R is $2\pi R$. Therefore

$$|f'(z)| \leq \frac{1}{2\pi} \left(\frac{M}{R^2} \right) (2\pi R) = \frac{M}{R}$$

This is true for every $R > 0$; now let $R \rightarrow \infty$ to find $f'(z) = 0$ for every $z \in \mathbb{C}$.

5.1 - Sequences and Series

Series: Basic Idea

- ▶ We know from real variable theory that many functions can be expressed as infinite series.
- ▶ For example,

$$\begin{aligned}e^x &= 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots \\&= \sum_{k=0}^{\infty} \frac{x^k}{k!}\end{aligned}$$

- ▶ To what extent does this theory extend to complex variables? Many of the definitions and theorems are similar...

Definitions

- **Definition:** A **series** is a sum

$$c_0 + c_1 + c_2 + \cdots = \sum_{k=0}^{\infty} c_k \text{ where the terms } c_k \in \mathbb{C}$$

- The n^{th} **partial sum** is $S_n = \sum_{k=0}^n c_k$.

- If $\lim_{n \rightarrow \infty} S_n$ exists and equals S (say), we say that $\sum_{k=0}^n c_k$

converges to S and we write $S = \sum_{k=0}^{\infty} c_k$

- If $\lim_{n \rightarrow \infty} S_n$ does not exist say the series **diverges**.

An Important Series

Theorem: Suppose $|z| < 1$. Then $\sum_{k=0}^{\infty} z^k = \frac{1}{1-z}$

Proof:

$$S_n = 1 + z + z^2 + \cdots + z^{n-1} + z^n$$

$$zS_n = z + z^2 + \cdots + z^{n-1} + z^n + z^{n+1}$$

$$S_n - zS_n = 1 - z^{n+1}$$

$$S_n(1 - z) = 1 - z^{n+1}$$

$$S_n = \frac{1 - z^{n+1}}{1 - z}$$

continued...

An Important Series, continued

Now let $n \rightarrow \infty$, so that $|z^{n+1}| \rightarrow 0$ since $|z| < 1$, leaving

$$\lim_{n \rightarrow \infty} S_n = \sum_{k=0}^{\infty} z^k = \frac{1}{1-z}$$

Convergence Tests

- ▶ Many convergence results for series of real terms extend to those with complex terms and the proofs are similar.
- ▶ **The Comparison Test:** Suppose $|c_k| \leq M_k$ for all $k \geq K$ (that is, eventually all of the c_k terms have modulus bounded by real numbers M_k .)

Then if $\sum_{k=0}^{\infty} M_k$ converges so does $\sum_{k=0}^{\infty} c_k$.

continued...

Convergence Tests, continued

- **The Ratio Test:** Suppose the series $\sum_{k=0}^{\infty} c_k$ is such that

$$\lim_{k \rightarrow \infty} \left| \frac{c_{k+1}}{c_k} \right| = L. \text{ Then}$$

- (i) If $L < 1$ the series converges
- (ii) If $L > 1$ the series diverges
- (iii) If $L = 1$ the test is inconclusive

Convergence Examples

Example: Determine the sum $\sum_{k=0}^{\infty} \frac{3}{(1+i)^k}$

Convergence Examples

Example: Determine whether $\sum_{k=0}^{\infty} \frac{k^2}{3^k}$ converges or diverges.

Convergence Examples

Example: Determine the largest open disk D on which

$$\sum_{k=0}^{\infty} \frac{(z-i)^k}{2^k} \text{ converges.}$$

Absolute Convergence

- ▶ **Definition:** The series $\sum_{k=0}^{\infty} c_k$ is called **absolutely convergent** if $\sum_{k=0}^{\infty} |c_k|$ converges.

- ▶ By the comparison test, taking $M_k = |c_k|$,

$$\sum_{k=0}^{\infty} |c_k| \text{ convergent} \implies \sum_{k=0}^{\infty} c_k \text{ convergent}$$

That is, absolute convergence implies convergence.

Pointwise Convergence

- ▶ Consider a function $F_n(z)$ defined on a set T , where $F_n(z)$ depends on both a non-negative integer n and $z \in \mathbb{C}$.

For example: $F_n(z) = \sum_{k=0}^n z^k = \frac{1 - z^{n+1}}{1 - z}$, and T is the disk $|z| < 1$.

- ▶ If for any $z \in \mathbb{C}$, $\lim_{n \rightarrow \infty} F_n(z)$ exists and equal $F(z)$, we say that F_n **converges pointwise** to F .
- ▶ **Definition:** F_n converges **pointwise** to F on T if for each $z \in T$, given $\epsilon > 0$ there is a natural number N (possibly depending on **both** ϵ and z) such that if $n > N$ then $|F_n(z) - F(z)| < \epsilon$.

Pointwise Convergence, Continued

- ▶ For $F_n(z) = \sum_{k=0}^n z^k = \frac{1 - z^{n+1}}{1 - z}$, we saw $F(z) = \frac{1}{1 - z}$, and again T is the disk $|z| < 1$.
- ▶ Notice: $|F_n(z) - F(z)| = \left| \frac{z^{n+1}}{1 - z} \right|$ depends on both n and z .
In order to make this difference small, n must be chosen with reference to the particular z being considered.
- ▶ Here $F_n(z) \rightarrow F(z)$ pointwise on T

Uniform Convergence

- ▶ Again consider a function $F_n(z)$ defined on a set T , where $F_n(z)$ depends on both a non-negative integer n and $z \in \mathbb{C}$.
- ▶ **Definition:** F_n converges **uniformly** to F on T if given $\epsilon > 0$ there is a natural number N (possibly depending on ϵ **but not on any particular z**) such that if $n > N$ then for any $z \in T$, $|F_n(z) - F(z)| < \epsilon$.
- ▶ Roughly speaking, if $F_n \rightarrow F$ uniformly, for n large enough the difference $|F_n(z) - F(z)|$ will be small for every $z \in T$.

Uniform Convergence, Continued

- ▶ Again consider $F_n(z) = \sum_{k=0}^n z^k = \frac{1 - z^{n+1}}{1 - z}$ and

$F(z) = \frac{1}{1 - z}$, but this time let T be the disk $|z| < 1/2$.

- ▶ Again

$$|F_n(z) - F(z)| = \left| \frac{z^{n+1}}{1 - z} \right| < \frac{(1/2)^{n+1}}{(1/2)} = \frac{1}{2^n}$$

- ▶ Notice: $|F_n(z) - F(z)|$ is bounded by an expression which is independent of z and which goes to zero as $n \rightarrow \infty$:
 $F_n \rightarrow F$ uniformly on T .

5.2 - Taylor Series

Taylor Series Definition

- ▶ **Definition:** Suppose f is analytic at z_0 . Then

$$\sum_{j=0}^{\infty} \frac{f^{(j)}(z_0)}{j!} (z - z_0)^j$$

is called the **Taylor Series** for f around z_0 .

- ▶ If $z_0 = 0$ the series above is instead called a **Maclaurin Series**

Taylor Series Example

Example: Construct the Maclaurin series for $f(z) = e^z$

Solution: $f(0) = f'(0) = f''(0) = f'''(0) = \dots = e^0 = 1$, so the Maclaurin series is

$$\sum_{j=0}^{\infty} \frac{f^{(j)}(0)}{j!} z^j = \sum_{j=0}^{\infty} \frac{1}{j!} z^j = 1 + z + \frac{z^2}{2!} + \frac{z^3}{3!} + \dots$$

The Main Result

- ▶ Under what conditions is a function **equal** to its Taylor series?
- ▶ **Theorem:** If f is analytic in a disk $D = \{|z - z_0| < R\}$, then

$$f(z) = \sum_{j=0}^{\infty} \frac{f^{(j)}(z_0)}{j!} (z - z_0)^j$$

for every z in D .

Furthermore, the series converges uniformly in any subdisk $D' = \{|z - z_0| \leq R' < R\}$.


- ▶ Consequently, the Taylor series will converge to $f(z)$ everywhere inside the largest disk centred at z_0 over which $f(z)$ is analytic.

Proof in the case $z_0 = 0$

Let C have radius $(R' + R)/2$.

For any z in D' ,

$$f(z) = \frac{1}{2\pi i} \int_C \frac{f(\zeta)}{\zeta - z} d\zeta, \text{ where}$$



$$= \frac{1}{2\pi i} \int_C f(\zeta) \left[\frac{1}{\zeta} \cdot \frac{1}{1 - z/\zeta} \right] d\zeta \quad \} \text{ notice } |z/\zeta| < 1$$

$$= \frac{1}{2\pi i} \int_C f(\zeta) \left[\frac{1}{\zeta} \cdot \left(\sum_{j=0}^n (z/\zeta)^j + \frac{(z/\zeta)^{n+1}}{1 - z/\zeta} \right) \right] d\zeta$$

Proof in the case $z_0 = 0$, continued

Splitting this last expression:

$$\begin{aligned} & \sum_{j=0}^n \frac{z^j}{j!} \left(\frac{j!}{2\pi i} \int_C \frac{f(\zeta)}{\zeta^{j+1}} d\zeta \right) + \frac{1}{2\pi i} \int_C \frac{f(\zeta)}{\zeta - z} \cdot \left(\frac{z}{\zeta} \right)^{n+1} d\zeta \\ &= \sum_{j=0}^n \frac{f^{(j)}(0)}{j!} z^j + \frac{1}{2\pi i} \int_C \frac{f(\zeta)}{\zeta - z} \cdot \left(\frac{z}{\zeta} \right)^{n+1} d\zeta \end{aligned}$$

Notice: as $n \rightarrow \infty$ the first sum becomes the desired Taylor series.

It remains to show that

$$\lim_{n \rightarrow \infty} \frac{1}{2\pi i} \int_C \frac{f(\zeta)}{\zeta - z} \cdot \left(\frac{z}{\zeta} \right)^{n+1} d\zeta = 0$$

Proof in the case $z_0 = 0$, continued

$$\frac{1}{2\pi i} \int_C \frac{f(\zeta)}{\zeta - z} \cdot \left(\frac{z}{\zeta}\right)^{n+1} d\zeta$$


On C ,

$$\left| \frac{1}{\zeta - z} \right| \leq \frac{1}{\left(\frac{R+R'}{2} - R'\right)} = \frac{2}{R - R'}$$

and

$$\left| \frac{z}{\zeta} \right|^{n+1} = \frac{|z|^{n+1}}{|\zeta|^{n+1}} \leq \left[\frac{R'}{\left(\frac{R'+R}{2}\right)} \right]^{n+1} = \left(\frac{2R'}{R' + R} \right)^{n+1} = \alpha^{n+1}$$

where $\alpha < 1$



Proof in the case $z_0 = 0$, continued

Using these bounds we have

$$\begin{aligned} & \left| \frac{1}{2\pi i} \int_C \frac{f(\zeta)}{\zeta - z} \cdot \left(\frac{z}{\zeta} \right)^{n+1} d\zeta \right| \\ & \leq \frac{1}{2\pi} \max_{\zeta \in C} |f(\zeta)| \left(\frac{2}{R - R'} \right) \alpha^{n+1} \\ & \rightarrow 0 \text{ as } n \rightarrow \infty \end{aligned}$$