
1. A methane molecule has a single carbon atom surrounded by four hydrogen atoms, like so:



The angles formed by the carbon-hydrogen bonds are all equal. Determine this common angle size.

- 2. A 100 lb weight is suspended from the ceiling by two ropes: one is a 3 ft rope which makes an angle α with the ceiling, the second a 4 ft rope which makes an angle β with the ceiling. The rope attachment points are 5 ft apart on the ceiling. Determine the angles α and β as well as the tension forces in the two ropes.
- 3. Give a parametrization of the line segment joining P(1, 2, 0) and Q(1, 3, -1).
- 4. What angle does the line of intersection of the planes 2x + y z = 0 and x + y + 2z = 0 make with the positive x-axis?
- 5. Let

$$f(x) = \begin{cases} \frac{\sin(x-y)}{|x|+|y|} & \text{if } |x|+|y| \neq 0\\ 0 & \text{if } (x,y) = (0,0) \end{cases}$$

Is *f* continuous at the origin?

- 6. Find parametric equations for the tangent line to the curve of intersection of the surfaces $x^2 + 2y + 2z = 4$ and y = 1 at the point (1, 1, 1/2).
- 7. Find the plane

$$\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$$

that passes through the point (2, 1, 2) and cuts off the least volume from the first octant.

- 8. Compute $\int_{0}^{8} \int_{\sqrt[3]{x}}^{2} \frac{1}{1+y^{4}} \, dy \, dx$
- 9. Draw the region in the xy-plane that lies inside the cardioid $r = 1 + \cos(\theta)$ but outside the circle r = 1. This region is the base of a cylinder which has top defined by the plane z = 7 x. Express the volume of the resulting solid as a triple integral in cylindrical coordinates.