

Question 1 [10]:

Determine $\int \frac{1}{x\sqrt{25-x^2}} dx$. (State your final answer without using inverse hyperbolic functions.)

Question 2:

(a)[4] Determine $\int \sin^3 x \cos^7 x \, dx$

(b)[6] Evaluate $\int_0^1 \arctan x \, dx$

Question 3 [10]:

Determine $\int \frac{3x^2 + 4}{x^3 + 2x} dx$.

Question 4:

(a)[5 points] Evaluate the improper integral $\int_3^5 \frac{1}{\sqrt{5-x}} dx$. Show all steps including any required limits.

(b)[5 points] Determine if $\int_1^\infty \frac{\sin^2 x}{x^2 + \sqrt{x}} dx$ converges or diverges. State reasons for your conclusion.


Question 5:

(a)[5 points] Use S_4 , Simpson's rule on four subintervals, to approximate $\int_0^4 \sqrt{3 + \cos^2(\pi x)} dx$.

(b)[5 points] The fourth derivative of $f(x) = \sqrt{3 + \cos^2(\pi x)}$ is between -300 and 160 for every x . If we wish to approximate $\int_0^4 \sqrt{3 + \cos^2(\pi x)} dx$ with accuracy 0.001 using Simpson's rule, how many subintervals are required?

Question 6:

- (a)[5 points] The base of a solid is the region bounded between the curve $y = \sqrt{1-x^2}$ and the x axis. Cross-sections perpendicular to both the base and the x -axis are squares. Find the volume of the solid.



- (b)[5 points] Find the area of the region bounded by the curves $4x + y^2 = 12$ and $y = x$.

