

Question 1: Solve the following system of equations **using either Gaussian or Gauss-Jordan elimination** (no credit will be given for using any other method). Use proper notation to state the row operations used at each step and clearly state the final solution.

$$\begin{aligned}y + 2z + 2w &= -6 \\x + 2y + 5z + 3w &= 2 \\2x + y + 5z + w &= -3\end{aligned}$$

Question 2: Each of the following matrices in reduced row echelon form (RREF) was obtained from a system of linear equations. For each of the reduced matrices,

- (a) If the solution is unique, write the solution.
- (b) If there is more than one solution, properly express the solution using parameters.
- (c) If there is no solution, then state "no solution"

(i) $\begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$ (the variables here are x, y, z .)

[1]

(ii) $\begin{bmatrix} 1 & 2 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$ (the variables here are x, y, z .)

[1]

(iii) $\begin{bmatrix} 1 & 0 & 4 & 5 & 1 \\ 0 & 1 & 1 & 3 & 2 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$ (the variables here are x, y, z, w .)

[1]

(iv) $\begin{bmatrix} 1 & 0 & 0 & 5 \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & 2 \end{bmatrix}$ (the variables here are x, y, z .)

[1]

Question 3: If the system

$$\begin{aligned} 2x - 4y &= 1 \\ ax + 3y &= 2 \end{aligned}$$

has no solution what must be the value of a ?

[2]

Question 4: For this problem use the following matrices:

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & 1 \\ 3 & 5 & 4 \\ 3 & 6 & 5 \end{bmatrix} \quad \mathbf{B} = \begin{bmatrix} 3 & -2 & 0 \\ -1 & -1 & 4 \end{bmatrix} \quad \mathbf{C} = \begin{bmatrix} 0 & 0 \\ 0 & 1 \\ 1 & 1 \end{bmatrix}$$

- (a) Compute $\mathbf{AB}^T - 2\mathbf{C}$ if it is defined.

[3]

- (b) Compute $\text{tr}((\mathbf{A} + \mathbf{I})(\mathbf{A} - \mathbf{I}))$ if it is defined.

[2]

- (c) Determine \mathbf{A}^{-1} if it exists.

[5]

Question 5: Determine the determinant of $\mathbf{A} = \begin{bmatrix} 1 & 1 & 1 & 1 & 1 \\ 1 & 2 & 3 & 4 & 5 \\ 1 & 1 & 4 & 3 & 2 \\ 1 & 1 & 1 & 3 & 4 \\ 1 & 1 & 1 & 1 & 5 \end{bmatrix}$.

(Row operations will make this much easier.)

[4]

Question 6: Use Cramer's rule to solve for y :

$$\begin{aligned}x + y - z &= 1 \\ 2x + 4y + 5z &= -2 \\ x + y + 2z &= -1\end{aligned}$$

Note that $\begin{vmatrix} 1 & 1 & -1 \\ 2 & 4 & 5 \\ 1 & 1 & 2 \end{vmatrix} = 6$.

[4]

Question 7: Suppose \mathbf{u} , \mathbf{v} and \mathbf{w} are any vectors in R^3 . Circle true (T) or false (F), as appropriate, for the following statements:

- | | | |
|--|---|---|
| (i) $\mathbf{u} \cdot (\mathbf{u} \times \mathbf{v}) = 0$ | T | F |
| (ii) $\mathbf{u} \times (\mathbf{u} \cdot \mathbf{v}) = 0$ | T | F |
| (iii) $\mathbf{u} \cdot \mathbf{u} = \ \mathbf{u}\ ^2$ | T | F |
| (iv) $\text{proj}_{\mathbf{u}} \mathbf{v} = \text{proj}_{\mathbf{v}} \mathbf{u}$ | T | F |
| (v) If \mathbf{u} , \mathbf{v} and \mathbf{w} are linearly independent, then $\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w}) = 0$ | T | F |

[5]

Question 8: Determine the equation of the line in R^3 that contains the point $(-1, 6, 0)$ and is orthogonal to the plane $4x + 2y - z = 5$. You may state your answer in either vector or parametric form.

[3]

Question 9: Find an equation of the plane passing through the points $P_1(-1, 0, 4)$, $P_2(-1, 4, 3)$ and $P_3(0, 6, -2)$. State your answer in the form

$$ax + by + cz = d$$

for appropriate constants a , b , c and d . (We saw three different methods for doing this problem; use whichever method you like.)

[5]

Question 10: Suppose $\mathbf{v}_1 = (1, a, 2)$, $\mathbf{v}_2 = (0, 2, a)$ and $\mathbf{v}_3 = (1, 1, 1)$ are linearly dependent. What are the possible values for a ?

[4]

Question 11: Let $\mathbf{u}_1 = (1, -1)$, $\mathbf{u}_2 = (1, 1)$ and $\mathbf{w} = (2, -3)$. Find the coordinate vector of \mathbf{w} relative to the basis $S = \{\mathbf{u}_1, \mathbf{u}_2\}$. That is, find $(\mathbf{w})_S$.

[4]

Question 12: Let $\mathbf{p}_1 = 1 + 2x + x^2$ and $\mathbf{p}_2 = 2 - x^2$. Find a third vector \mathbf{p}_3 so that $S = \{\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3\}$ is a basis for P_2 . (There are many possible correct answers here; give one answer with justification.)

[4]

Question 13: The matrix $\mathbf{A} = \begin{bmatrix} 1 & 2 & 0 & 0 & 3 \\ 1 & 2 & 1 & 0 & 2 \\ 3 & 6 & 1 & 0 & 8 \\ -1 & -2 & -2 & 0 & -1 \end{bmatrix}$ has RREF $\mathbf{R} = \begin{bmatrix} 1 & 2 & 0 & 0 & 3 \\ 0 & 0 & 1 & 0 & -1 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$.

(a) Give a basis and state the dimension of the row space of \mathbf{A} .

[2]

(b) Give a basis and state the dimension of the column space of \mathbf{A} .

[2]

(c) Give a basis and state the dimension of the null space of \mathbf{A} . (Equivalently, give a basis and state the dimension of the solution space of $\mathbf{Ax} = \mathbf{0}$.)

[3]

(d) Determine $\text{rank}(\mathbf{A})$ and $\text{nullity}(\mathbf{A})$.

[1]

(e) Determine $\text{rank}(\mathbf{A}^T)$ and $\text{nullity}(\mathbf{A}^T)$.

[1]

Question 14:

Let the transformation T_1 represent reflection about the yz -plane:

$$T_1 \left(\begin{bmatrix} a \\ b \\ c \end{bmatrix} \right) = \begin{bmatrix} -a \\ b \\ c \end{bmatrix} ,$$

and T_2 be the transformation which rotates a vector in R^3 counter-clockwise about the z -axis by an angle θ .

- (a) Determine the standard matrix **A** for the transformation T_1 .

[2]

- (b) Determine the standard matrix **B** for the transformation T_2 .

[3]

- (c) Determine the image of the vector $(1, 2, -1)$ if it is first reflected about the yz -plane and then rotated about the z -axis by $\pi/4$ (or 45° .)

[3]

- (d) For any vector (a, b, c) , will applying the transformation T_1 followed by T_2 produce the same result as first applying T_2 followed by T_1 ? Explain using matrices.

[2]

Question 15: Find the eigenvalues and corresponding eigenvectors (that is, the eigenpairs) for

$$\mathbf{A} = \begin{bmatrix} 2 & -1 \\ -2 & 3 \end{bmatrix}$$