

# Math 372 - Introductory Complex Variables

G.Pugh

Mar 29 2017

# Zeros and Singularities

# Zeros of Analytic Functions

- ▶ **Definition:** A **zero** of a function  $f$  is a point  $z_0$  where  $f$  is analytic and  $f(z_0) = 0$ .
- ▶ **Definition:**  $z_0$  is a **zero of order  $m$**  of  $f$  if  $f$  is analytic at  $z_0$  and  $f(z_0) = 0$ ,  $f'(z_0) = 0$ ,  $f''(z_0) = 0$ ,  $\dots$ ,  $f^{(m-1)}(z_0) = 0$ , but  $f^{(m)}(z_0) \neq 0$ .

# Zeros of Analytic Functions

- So, if  $f$  has a zero of order  $m$  at  $z_0$ , then the Taylor series for  $f$  about  $z_0$  takes the form

$$\begin{aligned}f(z) &= \frac{f^{(m)}(z_0)}{m!}(z - z_0)^m + \frac{f^{(m+1)}(z_0)}{(m+1)!}(z - z_0)^{m+1} + \dots \\&= (z - z_0)^m \left[ a_m + a_{m+1}(z - z_0) + a_{m+2}(z - z_0)^2 + \dots \right] \\&= (z - z_0)^m g(z)\end{aligned}$$

where  $g(z)$  is analytic at  $z_0$  and  $g(z_0) \neq 0$  in a neighbourhood of  $z_0$ .

- **Example:**  $f(z) = \cos(z) - 1 + z^2/2$  has a zero of order 4 at  $z = 0$  since

$$f(z) = \frac{z^4}{4!} - \frac{z^6}{6!} + \dots$$

# Isolated Singularities of Analytic Functions

- ▶ **Definition:** An **isolated singularity** of a function  $f$  is a point  $z_0$  such that  $f$  is analytic in some punctured disk  $0 < |z - z_0| < R$  but  $f$  is not analytic at  $z_0$  itself.
- ▶ **Example:**  $f(z) = \exp(z)/(z - i)$  has an isolated singularity at  $z = i$ .
- ▶ If  $f$  has an isolated singularity at  $z_0$ , then it has a Laurent series representation

$$f(z) = \sum_{j=-\infty}^{\infty} a_j(z - z_0)^j$$

in the punctured disk.

- ▶ Singularities are classified based on the form of the Laurent Series.

# Isolated Singularities of Analytic Functions

**Definition:** Suppose  $f$  has an isolated singularity at  $z_0$  and that

$$f(z) = \sum_{j=-\infty}^{\infty} a_j(z - z_0)^j$$

on  $0 < |z - z_0| < R$ .

- ▶ If  $a_j = 0$  for all  $j < 0$ , so that  $f(z) = \sum_{j=0}^{\infty} a_j(z - z_0)^j$ , then  $z_0$  is called a **removable singularity**.
- ▶ If  $a_{-m} \neq 0$  for some positive integer  $m$  but  $a_j = 0$  for all  $j < -m$ , then  $z_0$  is called a **pole of order  $m$**  of  $f$ .
- ▶ If  $a_j \neq 0$  for infinitely many  $j < 0$  then  $z_0$  is called an **essential singularity** of  $f$ .

# Removable Singularities

Suppose  $f$  has a removable singularity at  $z_0$ . Then

$$\begin{aligned} f(z) &= \sum_{j=0}^{\infty} a_j (z - z_0)^j \\ &= a_0 + a_1(z - z_0) + a_2(z - z_0)^2 + \dots \end{aligned}$$

**Example:**  $\frac{e^z - 1}{z} = 1 + \frac{z}{2!} + \frac{z^2}{3!} + \dots$

- ▶  $f$  is bounded in some punctured circular neighbourhood of  $z_0$
- ▶  $\lim_{z \rightarrow z_0} f(z)$  exists.
- ▶  $f$  can be redefined at  $z = z_0$  so that the new function is analytic at  $z_0$ . Define  $f(z_0) = a_0$ .

# Poles

Suppose  $f$  has a pole of order  $m$  at  $z_0$ . Then

$$f(z) = \frac{a_{-m}}{(z - z_0)^m} + \frac{a_{-m+1}}{(z - z_0)^{m-1}} + \cdots + a_0 + a_1(z - z_0) + a_2(z - z_0)^2 + \cdots$$

**Example:**  $\frac{\cos z}{z^2} = \frac{1}{z^2} - \frac{1}{2} + \frac{z^2}{4!} + \cdots$  has a pole of order 2 at  $z = 0$

- ▶  $(z - z_0)^m f(z)$  has a removable singularity at  $z_0$
- ▶  $\lim_{z \rightarrow z_0} |f(z)| = \infty$ .
- ▶ **Lemma:**  $f$  has a pole of order  $m$  at  $z_0$  if and only if  $f(z) = g(z)/(z - z_0)^m$  in some punctured neighbourhood of  $z_0$  where  $g$  is analytic and not zero at  $z_0$ .
- ▶ **Lemma:** If  $f$  has a zero of order  $m$  at  $z_0$  then  $1/f$  has a pole of order  $m$ . If  $f$  has a pole of order  $m$  at  $z_0$ , then  $1/f$  has a removable singularity at  $z_0$ , and  $1/f$  has a zero of order  $m$  at  $z_0$  if we define  $(1/f)(z_0) = 0$ .



# Essential Singularities

Suppose  $f$  has an essential singularity at  $z_0$ . Then

$$f(z) = \cdots + \frac{a_{-2}}{(z - z_0)^2} + \frac{a_{-1}}{(z - z_0)} + a_0 + a_1(z - z_0) + a_2(z - z_0)^2 + \cdots$$

**Example:**  $\exp(1/z) = 1 + \frac{1}{z} + \frac{1}{2!z^2} + \frac{1}{3!z^3} + \cdots$

**Theorem (*Picard*):** A function with an essential singularity at  $z_0$  assumes every complex number, with possibly one exception, as a value in any neighbourhood of  $z_0$ .

# Summary

**Theorem:** Suppose  $f$  has an isolated singularity at  $z_0$ . Then

- ▶  $z_0$  is a removable singularity  $\Leftrightarrow |f|$  is bounded near  $z_0 \Leftrightarrow \lim_{z \rightarrow z_0} f(z)$  exists  $\Leftrightarrow f$  can be redefined at  $z_0$  so that  $f$  is analytic at  $z_0$ .
- ▶  $z_0$  is a pole  $\Leftrightarrow \lim_{z \rightarrow z_0} |f(z)| = \infty \Leftrightarrow f(z) = g(z)/(z - z_0)^m$  in some punctured neighbourhood of  $z_0$  where  $g$  is analytic and not zero at  $z_0$ .
- ▶  $z_0$  is an essential singularity  $\Leftrightarrow |f(z)|$  is neither bounded near  $z_0$  nor goes to  $\infty$  as  $z \rightarrow z_0 \Leftrightarrow f$  assumes every complex number, with possibly one exception, as a value in any neighbourhood of  $z_0$ .

Can use this theorem to classify isolated singularities without constructing the Laurent Series.