

Question 1:

(a)[3 points] Let $f(x) = \sinh(\cosh^{-1} x)$. Find $f'(x)$.

$$f'(x) = \cosh(\cosh^{-1} x) \cdot \frac{1}{\sqrt{x^2 - 1}}$$

$$= \boxed{\frac{x}{\sqrt{x^2 - 1}}}$$

(b)[3 points] Find $f'(1)$ if $f(x) = \arctan(\sqrt{x})$.

$$f'(x) = \frac{1}{1 + (\sqrt{x})^2} \cdot \frac{1}{2} x^{-\frac{1}{2}}$$

$$f'(1) = \frac{1}{2} \cdot \frac{1}{2} \cdot 1 = \boxed{\frac{1}{4}}$$

(c)[4 points] Evaluate $\lim_{x \rightarrow \infty} x^2 \sin\left(\frac{1}{x^2}\right)$.

$$\lim_{x \rightarrow \infty} x^2 \sin\left(\frac{1}{x^2}\right) = \text{"}\infty \cdot 0\text{"} \quad \left. \vphantom{\lim_{x \rightarrow \infty} x^2 \sin\left(\frac{1}{x^2}\right)} \right\} \text{indeterminate form}$$

$$= \lim_{x \rightarrow \infty} \frac{\sin\left(\frac{1}{x^2}\right)}{x^{-2}} \sim \frac{0}{0} \quad \left. \vphantom{\lim_{x \rightarrow \infty} \frac{\sin\left(\frac{1}{x^2}\right)}{x^{-2}}} \right\} \text{indeterminate form}$$

$$\stackrel{H}{=} \lim_{x \rightarrow \infty} \frac{\cos\left(\frac{1}{x^2}\right) \cdot \cancel{(-2x^{-3})}}{\cancel{(-2x^{-3})}}$$

$$= \boxed{1}$$

Question 2:

(a) [3 points] Find $f(x)$ if $f'(x) = e^{2x} - \frac{1}{\sqrt{x}}$ and $f(0) = 1$.

$$f(x) = \int e^{2x} - x^{-\frac{1}{2}} dx = \frac{e^{2x}}{2} - 2x^{\frac{1}{2}} + C$$

$$f(0) = 1, \text{ so } \frac{e^{2 \cdot 0}}{2} - 2 \cdot 0^{\frac{1}{2}} + C = 1 \Rightarrow C = 1 - \frac{1}{2} = \frac{1}{2}$$

$$\therefore f(x) = \frac{e^{2x}}{2} - 2x^{\frac{1}{2}} + \frac{1}{2}$$

(b) [4 points] An object initially $s(0) = 2$ m above the surface of the moon is projected vertically upward with an initial velocity of $v(0) = 10$ m/s. Using the fact that acceleration due to gravity on the moon is $a(t) = -1.6$ m/s², derive the formula for $s(t)$, the height of the object above the moon's surface at time t seconds.

$$a(t) = -1.6$$

$$\therefore v(t) = \int -1.6 dt = -1.6t + C$$

$$v(0) = 10, \text{ so } -1.6 \cdot 0 + C = 10 \Rightarrow C = 10$$

$$\therefore v(t) = -1.6t + 10$$

$$\therefore s(t) = \int -1.6t + 10 dt = -\frac{1.6t^2}{2} + 10t + C$$

$$s(0) = 2, \text{ so } -\frac{1.6 \cdot 0^2}{2} + 10 \cdot 0 + C = 2 \Rightarrow C = 2$$

$$\therefore s(t) = -0.8t^2 + 10t + 2$$

(c) [3 points] Suppose $f(x)$ is a continuous function with the property that

$$\int_0^x f(t) dt = \sin(2x) - \int_0^x \cos(2t) f(t) dt.$$

Find a formula for $f(x)$. (Hint: differentiate both sides of the equation above.)

$$\frac{d}{dx} \left(\int_0^x f(t) dt \right) = \frac{d}{dx} \left(\sin(2x) - \int_0^x \cos(2t) f(t) dt \right)$$


$$f(x) = 2\cos(2x) - \cos(2x) f(x)$$

$$\therefore f(x)(1 + \cos(2x)) = 2\cos(2x)$$

$$f(x) = \frac{2\cos(2x)}{1 + \cos(2x)}$$

Question 3:

(a) [5 points] The following figure shows points on the graph of $y = f(x)$. Use the trapezoid rule to estimate $\int_0^7 f(x) dx$:



$$\begin{aligned} \int_0^7 f(x) dx &\approx \frac{\Delta x}{2} [f(0) + 2f(1) + 2f(2) + \dots + 2f(6) + f(7)] \\ &= \frac{1}{2} [3 + 2 \cdot 2 + 2 \cdot 2 + 2 \cdot 0 + 2 \cdot (-1) + 2 \cdot (-1) + 2 \cdot (-1) + 0] \\ &= \frac{1}{2} [3 + 4 + 4 + 0 + (-2) + (-2) + (-2) + 0] \\ &= \boxed{\frac{5}{2}} \end{aligned}$$

(b) [5 points] $f(x) = xe^x$ has second derivative $f''(x) = (2+x)e^x$. If the midpoint rule is being used to approximate $\int_0^2 xe^x dx$, how many subintervals are required in order to be accurate to within 0.01? (Recall, the error in using the midpoint rule to approximate $\int_a^b f(x) dx$ is at most $\frac{K(b-a)^3}{24n^2}$, where $|f''(x)| \leq K$ on $[a, b]$.)

$$|f''(x)| \leq |(2+x)e^x| \leq (2+2)e^2 = 4e^2 \text{ on } [0, 2],$$

so take $K = 4e^2$.

We want $\frac{K(b-a)^3}{24n^2} < 0.01$

$$\frac{4e^2(2-0)^3}{24n^2} < 0.01$$

$$\therefore \frac{4e^2}{(3)(0.01)} < n^2$$

$$\therefore 36.4 < n$$

$$\therefore \boxed{n = 32}$$

Question 4:

(a) [5 points] Evaluate $\int 3x^2 - x \sin(x^2) dx = I$

$$\begin{aligned} \therefore I &= 3 \int x^2 dx - \frac{1}{2} \int \sin(x^2) 2x dx \\ &= \frac{3x^3}{3} + \frac{1}{2} \cos(x^2) + C \\ &= \boxed{x^3 + \frac{1}{2} \cos(x^2) + C} \end{aligned}$$

(b) [5 points] Evaluate $\int_1^e x^{\frac{3}{2}} \ln x dx$.

$$\begin{aligned} \text{Let } I &= \int x^{\frac{3}{2}} \ln x dx \\ u &= \ln x \quad dv = x^{\frac{3}{2}} dx \\ du &= \frac{1}{x} dx \quad v = \frac{2}{5} x^{\frac{5}{2}} \end{aligned}$$

$$\begin{aligned} \therefore I &= \int u dv = uv - \int v du \\ &= \frac{2}{5} (\ln x) x^{\frac{5}{2}} - \int \frac{2}{5} x^{\frac{5}{2}} \frac{1}{x} dx \\ &= \frac{2}{5} (\ln x) x^{\frac{5}{2}} - \frac{2}{5} \int x^{\frac{3}{2}} dx \\ &= \frac{2}{5} (\ln x) x^{\frac{5}{2}} - \left(\frac{2}{5}\right)^2 x^{\frac{5}{2}} \end{aligned}$$

$$\begin{aligned} \int_1^e x^{\frac{3}{2}} \ln x dx &= \left[\frac{2}{5} (\ln x) x^{\frac{5}{2}} - \left(\frac{2}{5}\right)^2 x^{\frac{5}{2}} \right]_1^e \\ &= \left[\frac{2}{5} e^{\frac{5}{2}} - \left(\frac{2}{5}\right)^2 e^{\frac{5}{2}} \right] - \left[0 - \left(\frac{2}{5}\right)^2 \right] \\ &= \boxed{\frac{6}{25} e^{\frac{5}{2}} + \frac{4}{25}} \end{aligned}$$

Question 5:

(a) [5 points] Evaluate $\int \tan^2 x \sec^4 x dx = I$

$$I = \int \tan^2 x (1 + \tan^2 x) \sec^2 x dx$$

$$u = \tan x$$

$$du = \sec^2 x dx$$

$$\therefore I = \int u^2 (1 + u^2) du$$

$$= \int u^2 + u^4 du$$

$$= \frac{u^3}{3} + \frac{u^5}{5} + C$$

$$= \boxed{\frac{\tan^3 x}{3} + \frac{\tan^5 x}{5} + C}$$

(b) [5 points] Evaluate $\int \frac{\sqrt{x^2-1}}{x^3} dx$. (The identity $\sin(2\theta) = 2 \sin \theta \cos \theta$ may be useful here.)

$$x = \sec \theta$$

$$dx = \sec \theta \tan \theta d\theta$$

$$\therefore I = \int \frac{\sqrt{\sec^2 \theta - 1} \sec \theta \tan \theta d\theta}{\sec^3 \theta}$$

$$= \int \frac{\tan^2 \theta \sec \theta d\theta}{\sec^3 \theta}$$


$$= \int \sin^2 \theta d\theta$$

$$= \int \frac{1 - \cos(2\theta)}{2} d\theta$$

$$= \frac{\theta}{2} - \frac{1}{2} \frac{\sin(2\theta)}{2} + C$$

$$= \frac{\theta}{2} - \frac{1}{2} \sin \theta \cos \theta + C$$

$$\sec \theta = \frac{x}{1}, \text{ so}$$



$$\therefore I = \frac{\sec^{-1}(x)}{2} - \frac{1}{2} \frac{\sqrt{x^2-1}}{x} \frac{1}{x}$$

$$= \boxed{\frac{\sec^{-1}(x)}{2} - \frac{\sqrt{x^2-1}}{2x^2} + C}$$

Question 6:

(a) [5 points] Evaluate $\int \frac{1}{x^3 + 4x^2} dx = I$

$$\begin{aligned} \frac{1}{x^3 + 4x^2} &= \frac{1}{x^2(x+4)} = \frac{A}{x} + \frac{B}{x^2} + \frac{C}{x+4} \\ &= \frac{Ax(x+4) + B(x+4) + Cx^2}{x^2(x+4)} \\ &= \frac{(A+C)x^2 + (4A+B)x + 4B}{x^2(x+4)} \end{aligned}$$

$$\therefore \textcircled{1} A+C=0$$

$$\textcircled{2} 4A+B=0$$

$$\textcircled{3} 4B=1 \Rightarrow B=\frac{1}{4} \quad ; \quad \textcircled{2} \Rightarrow 4A+\frac{1}{4}=0, \therefore A=-\frac{1}{16}$$

$$\therefore \textcircled{1} \Rightarrow C=-A=\frac{1}{16}$$

$$\therefore I = \int \frac{(-1/16)}{x} + \frac{(1/4)}{x^2} + \frac{(1/16)}{x+4} dx$$

$$= \boxed{-\frac{1}{16} \ln|x| + \frac{1}{4} \frac{1}{x} + \frac{1}{16} \ln|x+4| + C}$$

(b) [5 points] Evaluate the improper integral $\int_0^3 \frac{x}{\sqrt{9-x^2}} dx$

$$\int_0^3 \frac{x}{\sqrt{9-x^2}} dx = \lim_{b \rightarrow 3^-} \int_0^b \frac{x}{\sqrt{9-x^2}} dx$$


$$= \lim_{b \rightarrow 3^-} \left[-\sqrt{9-x^2} \right]_0^b$$

$$= \lim_{b \rightarrow 3^-} \left(-\sqrt{9-b^2} + \sqrt{9-0^2} \right)$$


$$= \boxed{3}$$

Question 7:

(a) [5 points] Find the area of the shaded region:




$$\begin{aligned}
 A &= \int_0^{\pi/2} \sin x - \frac{2}{\pi} x \, dx + \int_{\pi/2}^{\pi} \left(\frac{2}{\pi}\right) x - \sin x \, dx \\
 &= \left[-\cos x - \frac{1}{\pi} x^2\right]_0^{\pi/2} + \left[\frac{1}{\pi} x^2 + \cos x\right]_{\pi/2}^{\pi} \\
 &= \left[(-\cos(\frac{\pi}{2}) - \frac{1}{\pi}(\frac{\pi}{2})^2) - (-\cos(0) - 0)\right] + \left[\left(\frac{1}{\pi} \pi^2 + \cos(\pi)\right) - \left(\frac{1}{\pi}(\frac{\pi}{2})^2 + \cos(\frac{\pi}{2})\right)\right] \\
 &= -\frac{\pi}{4} + 1 + \pi - 1 - \frac{\pi}{4} = \boxed{\frac{\pi}{2}}
 \end{aligned}$$

(b) [5 points] The shaded region is rotated about the vertical line $x = \pi/2$; set up the integral representing the volume of the resulting solid. DO NOT EVALUATE THE INTEGRAL.

$$\therefore V = \int_0^{\pi/2} 2\pi \left(\frac{\pi}{2} - x\right) \left(\sin(x) - \frac{2}{\pi} x\right) dx$$

Question 8: Consider the graph of the following ellipse:



If the ellipse is rotated about the x -axis the resulting solid is called an ellipsoid (which looks rather like a watermelon).

(a) [3 points] Isolate y in the equation above to find a function which describes the top half of the ellipse.

$$\frac{x^2}{16} + \frac{y^2}{4} = 1$$

$$\therefore y = \sqrt{4\left(1 - \frac{x^2}{16}\right)}$$

(b) [7 points] Use your result in (a) to find the volume of the ellipsoid.

Using disks:

$$V = \int_{-4}^4 \pi \left(\sqrt{4\left(1 - \frac{x^2}{16}\right)} \right)^2 dx$$


$$= 2\pi \int_0^4 4\left(1 - \frac{x^2}{16}\right) dx$$

$$= 8\pi \left[x - \frac{x^3}{48} \right]_0^4$$

$$= 8\pi \left[4 - \frac{64}{48} \right]$$

$$= \boxed{\frac{64\pi}{3}}$$

Question 9: A rectangular fish tank of length 4 m, width 2 m and height 3 m contains water to a depth of 2 m. Recall that the density of water is $\rho = 1000 \text{ kg/m}^3$ and acceleration due to gravity is $g = 9.8 \text{ m/s}^2$.



(a) [5 points] How much work is required to pump all of the water out over the edge of the tank?

Weight of slice of water a depth $x = (4)(2)\Delta x \rho g$

\therefore Work required to lift this slice to top of tank is $(4)(2)\Delta x \rho g x = 8\rho g x \Delta x$

\therefore Total work $W = \int_{x=1}^{x=3} 8\rho g x \, dx$

$$= 8\rho g \left[\frac{x^2}{2} \right]_1^3$$

$$= 8\rho g \frac{9-1}{2}$$

$$= 32\rho g$$

$$= (32)(1000)(9.8)$$

$$= \boxed{313,600 \text{ N}\cdot\text{m}}$$

(b) [5 points] What is the hydrostatic force (force due to water pressure) exerted on one of the long sides of the tank? Recall that pressure P as a function of depth h is $P(h) = \rho gh$ where ρ is the density of the liquid and g is acceleration due to gravity.

area of strip of width Δx is $4\Delta x$

\therefore force on this strip is $4\Delta x \rho g (x-1)$

\therefore Total force $F = \int_{x=1}^{x=3} 4\rho g (x-1) \, dx$

$$= 4\rho g \int_1^3 (x-1) \, dx$$

$$= 4\rho g \left[\frac{x^2}{2} - x \right]_1^3$$

$$= 4\rho g \left[\frac{3}{2} + \frac{1}{2} \right]$$

$$= 8\rho g = (8)(1000)(9.8) = \boxed{78400 \text{ N}}$$

Question 10: The fish population in a large lake is infected by a disease at time $t = 0$, and the declining fish population is described by the differential equation

$$\frac{dP}{dt} = -k\sqrt{P}.$$

Here k is a positive constant and $P(t)$ is the fish population at time t weeks. Suppose there were initially 90,000 fish in the lake and that 40,000 remain after 6 weeks.

(a)[7 points] Solve the differential equation to find a formula for $P(t)$.

$$\int \frac{1}{\sqrt{P}} dP = \int -k dt$$

$$2\sqrt{P} = -kt + C$$

$$P(0) = 90,000, \text{ so}$$

$$2\sqrt{90,000} = -k \cdot 0 + C$$

$$\therefore C = 600$$

$$\therefore 2\sqrt{P} = -kt + 600$$

$$P(6) = 40,000, \text{ so}$$

$$2\sqrt{40,000} = -k \cdot 6 + 600$$

$$\therefore k = \frac{600 - 2 \cdot 200}{6}$$

$$= \frac{100}{3}$$

$$\therefore 2\sqrt{P} = 600 - \frac{100}{3}t$$

$$P(t) = \left(300 - \frac{50}{3}t\right)^2$$

(b)[3 points] Use your result in (a) to find the time required for the fish population to reduce to 10,000.

$$\text{Solve } \left(300 - \frac{50}{3}t\right)^2 = 10,000$$

$$300 - \frac{50}{3}t = 100$$

$$200 = \frac{50}{3}t$$

$$t = \frac{3 \cdot 200}{50} = \boxed{12 \text{ weeks}}$$

Question 11: Recall that $\sinh(x) = \frac{e^x - e^{-x}}{2}$, and that the Maclaurin series for e^x is

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots$$

(a)[4 points] Find the first three non-zero terms of the Maclaurin series for $f(x) = \sinh(x^2)$.

$$\begin{aligned} \sinh(x^2) &= \frac{e^{x^2} - e^{-x^2}}{2} \\ &= \frac{\left(1 + x^2 + \frac{x^4}{2!} + \frac{x^6}{3!} + \dots\right) - \left(1 - x^2 + \frac{x^4}{2!} - \frac{x^6}{3!} + \dots\right)}{2} \\ &= \boxed{x^2 + \frac{x^6}{3!} + \frac{x^{10}}{5!} + \dots} \end{aligned}$$

(b)[3 points] Use a Maclaurin series to evaluate the limit

$$\begin{aligned} &\lim_{x \rightarrow 0} \frac{e^x - 1 - x - (x^2/2)}{x^3} \\ &= \lim_{x \rightarrow 0} \frac{\left(1 + x + \frac{x^2}{2} + \frac{x^3}{3!} + \dots\right) - 1 - x - \frac{x^2}{2}}{x^3} \\ &= \lim_{x \rightarrow 0} \frac{1}{3!} + \frac{x^2}{4!} + \dots \\ &= \boxed{\frac{1}{6}} \end{aligned}$$

(c)[3 points] Suppose $f(x)$ is a function such that $f(2) = 3$, $f'(2) = 0$, $f''(2) = -1$ and $f'''(2) = 2$. Use a Taylor polynomial of degree 3 to approximate $f(2.1)$. Round your final answer to three decimals.

$$\begin{aligned} T_3(x) &= f(2) + f'(2)(x-2) + \frac{f''(2)}{2}(x-2)^2 + \frac{f'''(2)}{3!}(x-2)^3 \\ &= 3 - \frac{1}{2}(x-2)^2 + \frac{2}{3!}(x-2)^3 \end{aligned}$$

$$\begin{aligned} \therefore f(2.1) &\approx T_3(2.1) = 3 - \frac{1}{2}(0.1)^2 + \frac{2}{3!}(0.1)^3 \\ &= \boxed{2.995} \end{aligned}$$