



Question 1:


(a)[4] Find all values of x in $[0, 2\pi]$ that satisfy the equation $\sqrt{2} \sin x - 1 = 0$.



(b)[3] Determine the exact value of $\cos(4\pi/3)$.



(c)[3] Determine $\tan \theta$ if $\cos \theta = 2/5$ and $0 < \theta < \pi/2$.



Question 2:

Evaluate the following limits, if they exist:

$$(a)[3] \quad \lim_{x \rightarrow -1} \frac{x^2 + 4x + 3}{x + 1} = \lim_{x \rightarrow -1} \frac{\cancel{(x+1)}(x+3)}{\cancel{(x+1)}}$$

$$= \boxed{2}$$

$$(b)[4] \quad \lim_{x \rightarrow 1} \frac{x-1}{\sqrt{x+8}-3} \cdot \frac{\sqrt{x+8}+3}{\sqrt{x+8}+3}$$

$$= \lim_{x \rightarrow 1} \frac{\cancel{(x-1)}(\sqrt{x+8}+3)}{\cancel{(x-1)}}$$

$$= \boxed{6}$$

$$(c)[3] \quad \lim_{x \rightarrow 0} \frac{\sqrt{5x+9}-3}{x-3} = \frac{0}{-3} = \boxed{0} \quad (\text{Direct Sub.})$$

Question 3: For this question use the functions $f(x) = \sqrt{x+1}$, $g(x) = \frac{1}{x^2-5}$, $h(x) = \frac{1}{x}$.

(a)[3] Determine $(h \circ g \circ f)(x)$ and simplify.

$$\begin{aligned} (h \circ g \circ f)(x) &= h(g(f(x))) \\ &= \frac{1}{\left[\frac{1}{(\sqrt{x+1})^2-5}\right]} \\ &= \boxed{x-4} \end{aligned}$$

(b)[3] Determine the domain of $(g \circ f)(x)$.

$$(g \circ f)(x) = g(f(x)) = \frac{1}{(\sqrt{x+1})^2-5} = \frac{1}{x-4}$$

Require: $x+1 \geq 0 \Rightarrow x \geq -1$
and $x \neq 4$

\therefore domain is

$$\boxed{[-1, 4) \cup (4, \infty)}$$

(c)[2] Find a function $q(x)$ so that $g(x) = (q \circ h)(x)$.

Want $q(h(x)) = \frac{1}{x^2-5}$

$$\therefore q(x) = \frac{1}{\left(\frac{1}{x}\right)^2-5}$$

(d)[2] Find a function $p(x)$ so that $g(x) = (h \circ p)(x)$.

Want $h(p(x)) = \frac{1}{x^2-5}$

$$\therefore p(x) = x^2-5$$

Question 4:

(a)[5] Determine $\lim_{x \rightarrow 0} x^4 \cos(\sqrt{1+x^2})$.

(State any theorems used, like the the Squeeze Theorem, for example, and be sure to state the conditions necessary to justify use of the theorem.)

$$-x^4 \leq x^4 \cos(\sqrt{1+x^2}) \leq x^4$$

$$\lim_{x \rightarrow 0} -x^4 = 0 = \lim_{x \rightarrow 0} x^4$$

\therefore by the Squeeze Theorem, $\lim_{x \rightarrow 0} x^4 \cos(\sqrt{1+x^2}) = 0$.

Or, simply apply the direct substitution property:

$$\lim_{x \rightarrow 0} x^4 \cos(\sqrt{1+x^2}) = 0^4 \cdot \cos(\sqrt{1+0}) = 0$$

(b)[5] Evaluate the following limit if it exists: $\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta + \tan \theta}$

$$\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta + \frac{\sin \theta}{\cos \theta}} \quad \frac{0}{0}$$

$$= \lim_{\theta \rightarrow 0} \frac{\left(\frac{\sin \theta}{\theta}\right)}{\frac{\theta}{\theta} + \frac{\sin \theta}{\theta} \cdot \frac{1}{\cos \theta}}$$

$$= \frac{1}{1+1 \cdot 1}$$

$$= \boxed{\frac{1}{2}}$$

Question 5: Evaluate the following limits, if they exist:

$$(a)[2] \quad \lim_{x \rightarrow 2} \frac{\cos(\pi x)}{(x-2)^2} \quad \left. \begin{array}{l} \} \rightarrow "1" \\ \} \rightarrow 0 \end{array} \right\}$$

∴ limit does not exist.

$$(b)[4] \quad \lim_{x \rightarrow 0^-} \left(\frac{1}{x} + \frac{1}{|x|} \right) = \lim_{x \rightarrow 0^-} \frac{1}{x} - \frac{1}{x} \quad \text{since } x < 0$$

$$= \lim_{x \rightarrow 0^-} 0$$

$$= \boxed{0}$$

$$(c)[4] \quad \lim_{x \rightarrow 0} \frac{\left[\frac{1}{x-1} + \frac{1}{x+1} \right]}{x}$$

$$= \lim_{x \rightarrow 0} \frac{\left(\frac{x+1 + x-1}{(x-1)(x+1)} \right)}{\left(\frac{x}{1} \right)}$$

$$= \lim_{x \rightarrow 0} \frac{2x}{(x-1)(x+1)} \cdot \frac{1}{x}$$

$$= \boxed{-2}$$