

Supplemental Notes on Series

A brief introduction to Taylor and power series

VIU MATH 122 - Summer 2017

1 Introduction

These notes give a brief introduction to two very important and closely related calculus topics: the approximation of functions using certain types of polynomials, and the representation of functions using infinite series (that is, the sum of an infinite number of terms.)

For example, in a precalculus course you may have seen the *geometric series*

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + x^4 + \dots$$

which is true for any value of x between -1 and 1 . **By writing equality here we mean that the difference between $1/(1-x)$ and $1 + x + x^2 + x^3 + x^4 + \dots + x^n$ approaches zero as n grows to infinity.** In this way, for each natural number n the function $f(x) = 1/(1-x)$ is well approximated by the polynomial $p(x) = 1 + x + x^2 + x^3 + x^4 + \dots + x^n$. As n grows to infinity we get true equality of $f(x) = 1/(1-x)$ with the infinite series $1 + x + x^2 + x^3 + x^4 + \dots$.

Many other functions have such representations. For example, the familiar exponential function has representation

$$e^x = 1 + x + \frac{x^2}{2} + \frac{x^3}{2 \cdot 3} + \frac{x^4}{2 \cdot 3 \cdot 4} + \dots$$

while for the sine function,

$$\sin x = x - \frac{x^3}{2 \cdot 3} + \frac{x^5}{2 \cdot 3 \cdot 4 \cdot 5} - \dots$$

These last two series are valid for all real numbers x . In these notes we make these notions precise by looking at the errors inherent in the polynomial approximations, and the conditions under which a given function has an infinite series representation.

These infinite series representations are important for a number of reasons. One is that they form the basis of the methods used to evaluate functions. At the most fundamental level, numerical calculation (by human or machine) involves only the operations of addition, subtraction, multiplication and division, the same four operations required to add up the terms in a series. If, for example, $\sin(37^\circ)$ is required in a calculation, how do we (or our calculators) come up with the decimal approximation 0.60181502315? The procedure used is based on the series representation of $\sin x$ above.

Another important reason for the study of series is found in the solution of real-world physics and engineering problems. Often the solutions to such problems cannot be expressed as finite combinations of elementary functions (polynomial, trigonometric, exponential, etc) and an infinite series is the only known solution representation.

2 Some Definitions

To describe our results we need a few definitions. These are likely already familiar to you but we include them here for review and reference.

Definition 2.1 An **interval** is a set of numbers which form a segment of the real number line. For example, $(0, 1)$, $[0, 1]$, $(-3, \infty)$, $(-\infty, \pi]$, $(-\infty, \infty)$ are all intervals.

An **open interval** is an interval which includes neither its upper nor lower boundaries and so uses only "(" and ")" in its representation. Examples of open intervals are

$$(0, 1), (-2, \infty) \text{ and } (-\infty, \infty) .$$

A **closed interval** is an interval which includes both its upper and lower boundaries, such as $[0, 1]$ or $[e, \pi]$.

Intervals which include one boundary but not the other are sometimes call **half open**, such as $(-\infty, 1]$ or $[0, 7)$.

The letter \mathcal{I} is often used to denote a general interval.

Definition 2.2 A function f is **differentiable** at a point $x = a$ if $f'(a)$ exists, meaning that the limit defining the derivative

$$\lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$$

exists.

A function is **n-times differentiable** at a point $x = a$ if the n^{th} derivative $f^{(n)}(a)$ exists (and consequently $f'(a)$, $f''(a)$, \dots , $f^{(n)}(a)$ all exist.)

A function is **differentiable on an open interval** \mathcal{I} if it is differentiable at every point x in \mathcal{I} , and similarly for **n-times differentiable on an open interval**.


For example, $f(x) = x^{7/3}$ is twice differentiable on $(-1, 1)$ since $f''(x) = (28/9)x^{1/3}$ exists for every x in $(-1, 1)$. However, $f(x) = x^{7/3}$ is not 3-times differentiable on $(-1, 1)$ since at $x = 0$

$$\begin{aligned} f'''(0) &= \lim_{h \rightarrow 0} \frac{f''(0+h) - f''(0)}{h} \\ &= \lim_{h \rightarrow 0} \frac{(28/9)h^{1/3} - 0}{h} \\ &= \lim_{h \rightarrow 0} (28/9)h^{-2/3} \end{aligned}$$

which fails to exist.

3 Linear Approximation

Before looking at infinite series, we first develop a class of polynomials useful in the approximation of functions. The simplest of these approximating polynomials is something you have seen already when

Figure 1: $f(x)$, $T_1(x)$

you studied linear approximation, so we'll begin by reviewing that topic and then extend it with some new theory.

Recall the **linear approximation** (or **tangent line approximation**) to a function f at a point $x = a$: If f is differentiable at a , then for x near a

$$f(x) \approx f(a) + f'(a)(x - a) .$$

Here $y = f(a) + f'(a)(x - a)$ is just the equation of the tangent line to $y = f(x)$ at the point where $x = a$, so linear approximation is simply the idea of using the tangent line to approximate function values.

Letting $T_1(x) = f(a) + f'(a)(x - a)$, the function describing the tangent line, notice that

$$T_1(a) = f(a) \text{ and } T_1'(a) = f'(a) .$$

That is, the relatively simple function $T_1(x)$ agrees with $f(x)$ in function value and first derivative at $x = a$. The function $T_1(x)$ is called the **Taylor polynomial of degree 1 for f at a** . We see from plots of $y = f(x)$ and $y = T_1(x)$ in Figure 1 that for x near a the vertical distance between points $(x, f(x))$ and $(x, T_1(x))$ is small, but that this distance increases as x moves away from a .

We would like to say something more precise about how good the approximation $f(x) \approx T_1(x)$ is. Let $R_1(x)$ be the difference between $f(x)$ and $T_1(x)$:

$$R_1(x) = f(x) - [f(a) + f'(a)(x - a)] .$$

Think of $R_1(x)$ as the error in the approximation $f(x) \approx f(a) + f'(a)(x - a)$. How large can this error possibly be? The answer to this question, called a **bound** on the approximation error, can be found using the following theorem, the proof of which is in the Appendix:

Theorem 3.1 Suppose that f is twice differentiable on an open interval \mathcal{I} containing containing a . Then for each x in \mathcal{I}

$$R_1(x) = \frac{f''(z)}{2}(x - a)^2$$

for some z between a and x .

Notice here that z is some unknown number between a and x , so we cannot in general give the exact value of $R_1(x)$. However, by analyzing f'' we can establish an **upper bound** on $|f''(z)|$ (that is, a greatest possible size), and thereby give a bound on the error term $|R_1(x)|$ itself. This is illustrated in the following example.

Example 3.1

Let $f(x) = x + \ln x$.

- (i) Find the linear approximation to f at $a = 1$.
- (ii) Use the linear approximation to approximate $f(1.1)$.
- (iii) Give an error bound for your approximation.

Solution

- (i) Here

$$f(x) = x + \ln x \quad \text{and} \quad f'(x) = 1 + \frac{1}{x}.$$

Evaluating these at $a = 1$ gives

$$f(1) = 1 \quad \text{and} \quad f'(1) = 2,$$

so we have

$$\begin{aligned} T_1(x) &= f(a) + f'(a)(x - a) \\ &= 1 + 2(x - 1) \end{aligned}$$

- (ii) The linear approximation is $f(1.1) \approx T_1(1.1) = 1 + 2(1.1 - 1) = 1.2$

- (iii) We require an upper bound on $|R_1(1.1)|$. By Theorem 3.1

$$R_1(1.1) = \frac{f''(z)}{2}(1.1 - 1)^2$$

for some $1 < z < 1.1$. Here $f''(z) = -1/z^2$, so

$$|R_1(1.1)| = \left| \frac{-1}{2z^2}(1.1 - 1)^2 \right|.$$

Now ask: what is the largest possible size of $|R_1(1.1)|$ for $1 < z < 1.1$? In this case, allowing z to approach 1 makes $|R_1(1.1)|$ as large as possible, so setting $z = 1$ we have

$$|R_1(1.1)| \leq \left| \frac{-1}{2(1)^2}(1.1 - 1)^2 \right| = \frac{1}{200}.$$

That is,

$$|f(1.1) - T_1(1.1)| \leq \frac{1}{200}.$$

Note the following points about the error bound determination in this example:

1. When bounding the error in the approximation, it is the absolute value of the error term which is of interest.
2. Theorem 3.1 guarantees that the exact value of $R_1(1.1)$ occurs at some z in the open interval $(1, 1.1)$. However, the upper bound on $|R_1(1.1)|$ over the interval $(1, 1.1)$ may occur as $z \rightarrow 1$ or $z \rightarrow 1.1$, as was the case here. When determining error bounds, always consider the limiting value of the error as z approaches the endpoints of the interval.
3. In this particular example $|R_1(1.1)|$ is a decreasing function of z , so the bound on $|R_1(1.1)|$ occurred at one of the endpoints of the interval $(1, 1.1)$. This is not uncommon, but needn't always be the case: it depends on the form of the error term. Finding a bound on $|R_1(x)|$ that is best possible may sometimes require the optimization techniques studied in differential calculus.

□

The previous example shows that $T(1.1)$ is within $1/200$ or 0.005 of the true value of $f(1.1)$. A calculator check shows that this is indeed the case:

$$|f(1.1) - T(1.1)| = |(1.1 + \ln(1.1)) - (1 + 2(1.1 - 1))| \doteq 0.0047,$$

a remarkably accurate result given the simple form of the approximating function $T_1(x)$. In the sections to follow we will see how to improve the accuracy even further by extending $T_1(x)$ to a more general polynomial.

Section 3 Exercises

- 3.1** Determine the linear approximation $T_1(x)$ for $f(x) = e^{-x} + e^{-2x}$ at $a = 0$. $e^{-0.1}$. Give an error bound for your approximation.
- 3.2** Determine the linear approximation $T_1(x)$ for $f(x) = \sqrt{x}$ at $a = 1$ and use it to approximate $\sqrt{1.05}$. Give an error bound for your approximation.
- 3.3** Determine the linear approximation $T_1(x)$ for $f(x) = e^x$ at $a = 0$ and use it to approximate
- 3.4** Determine $T_1(x)$ for $f(x) = 1/(x + 2)$ at $a = 2$ and use it to approximate $f(3/2)$. Give an error bound for your approximation.
- 3.5** Use a linear approximation $T_1(x)$ for $f(x) = x/(x + 1)$ to approximate $f(1.3)$. Give an error bound for your approximation.

3.6 Determine $T_1(x)$ for $f(x) = \sqrt{x^2 + 9}$ at $a = -4$ and use it to approximate $f(-3.9)$. Give an error

bound for your approximation.

Answers

3.1 $T_1(x) = 2 - 3x$

3.2 $T_1(x) = 1 + \frac{1}{2}(x - 1); \sqrt{1.05} \approx T_1(1.05) = \frac{41}{40};$
 $|R_1(1.05)| \leq \frac{1}{3200}$

3.3 $T_1(x) = 1 + x; e^{-0.1} \approx T_1(-0.1) = \frac{9}{10};$
 $|R_1(-0.1)| \leq \frac{1}{200}$

3.4 $T_1(x) = \frac{1}{4} - \frac{1}{16}(x - 2); f(3/2) \approx T_1(3/2) = \frac{9}{32}; |R_1(3/2)| \leq \frac{2}{7^3}$

3.5 $T_1(x) = \frac{x+1}{4}; f(1.3) \approx T_1(1.3) = \frac{23}{40}; |R_1(1.3)| \leq \frac{9}{800}$

3.6 $T_1(x) = 5 - \frac{4(x+4)}{5}; f(-3.9) \approx T_1(-3.9) = \frac{123}{25}$
 $|R_1(-3.9)| \leq \frac{9}{200[(3.9)^2 + 9]^{3/2}} \approx 0.0004$

4 Taylor & Maclaurin Polynomials

We saw in the previous section that $T_1(x)$ does a reasonable job of approximating $f(x)$ for x near a . That is, the graph of $y = T_1(x)$ (a line) approximates that of $y = f(x)$ near $x = a$. Let's now improve the approximation by using polynomials of degree greater than 1, the graphs of which are **curves** and so one would expect that they would better approximate the graph of $y = f(x)$ near $x = a$.

Let n be a positive integer and suppose that f is n -times differentiable at $x = a$. Let's now generalize the linear approximation idea by finding a polynomial $T_n(x)$ of degree n which agrees with $f(x)$ in function value and first n -derivatives at $x = a$. That is, let's find $T_n(x)$ so that

$$\begin{aligned} T_n(a) &= f(a), \\ T'_n(a) &= f'(a), \\ T''_n(a) &= f''(a), \\ &\vdots \\ T_n^{(n)}(a) &= f^{(n)}(a) \end{aligned}$$

Since T_n has degree n it can be expressed in the general form

$$T_n(x) = b_0 + b_1(x - a) + b_2(x - a)^2 + \cdots + b_n(x - a)^n$$

for some constants b_0, b_1, \dots, b_n . To determine the constants b_0, b_1, \dots, b_n proceed as follows:

► Since $T_n(a) = f(a)$ we must have

$$f(a) = b_0 + b_1(a - a) + b_2(a - a)^2 + \cdots + b_n(a - a)^n$$

so that $b_0 = f(a)$.

► Since $T'_n(a) = f'(a)$ we must have

$$f'(a) = b_1 + 2b_2(a - a) + \cdots + nb_n(a - a)^{n-1}$$

so that $b_1 = f'(a)$.

► Since $T_n''(a) = f''(a)$ we must have

$$f''(a) = 2b_2 + (3)(2)b_3(a - a) + \cdots + n(n - 1)b_n(a - a)^{n-2}$$

so that $b_2 = \frac{f''(a)}{2}$.

► Continuing in this way we find in general

$$b_k = \frac{f^{(k)}(a)}{k!}, k = 0, 1, \dots, n$$

We can now define the **Taylor polynomial** of degree n for f at a :

Definition 4.1 If f is n -times differentiable at a , the **Taylor polynomial** of degree n for f at a is

$$T_n(x) = f(a) + f'(a)(x - a) + \frac{f''(a)}{2}(x - a)^2 + \frac{f'''(a)}{3!}(x - a)^3 + \cdots + \frac{f^{(n)}(a)}{n!}(x - a)^n$$

If $a = 0$, $T_n(x)$ is instead called a **Maclaurin polynomial** of degree n for f .

Some comments about this definition:


(i) Here '!' is **factorial notation**. $n!$ is read " n factorial", so $5!$ is read "five-factorial". The factorial function is defined on the non-negative integers as follows:

$$\begin{aligned} 0! &= 1 \\ 1! &= 1 \\ 2! &= (1)(2) \\ 3! &= (1)(2)(3) \\ &\vdots \\ k! &= (1)(2) \cdots (k) \end{aligned}$$

This can also be defined recursively as $0! = 1$ and $k! = k(k - 1)!$ for $k \geq 1$.

(ii) Notice that the first two terms of $T_n(x)$ give $T_1(x)$, the first three terms give $T_2(x)$, and so on. In other words, once you find $T_n(x)$ you also have $T_k(x)$ for $k = 1, 2, \dots, n - 1$.

Since $T_n(x)$ agrees with $f(x)$ in function value and first n -derivatives at $x = a$ we expect $f(x) \approx T_n(x)$ for x near a . See Figure 2 for an illustration of the improved approximations given by $T_1(x)$, $T_2(x)$ and $T_5(x)$.

Figure 2: $f(x)$, $T_1(x)$, $T_2(x)$, $T_5(x)$

Taylor's Theorem

As with linear approximation, when using a Taylor polynomial to approximate a function we would like an estimate of the error involved. There are several methods for doing this, one of which is a generalization of the error estimate we found for linear approximation. To state the result let

$$R_n(x) = f(x) - T_n(x)$$

be the error in the approximation.

Theorem 4.1 Suppose f is $(n + 1)$ -times differentiable on an open interval \mathcal{I} containing containing a . Then for each x in \mathcal{I}

$$R_n(x) = \frac{f^{(n+1)}(z)}{(n+1)!} (x - a)^{n+1}$$

for some z between a and x .

The proof, given in the Appendix, is nearly identical to the $n = 1$ case we saw before.

Putting all this together we arrive at the main result of this section: the formal statement of Taylor's Theorem, also known as **Taylor's Formula**:

Taylor's Theorem

Suppose f is $(n+1)$ -times differentiable on an open interval \mathcal{I} containing a . Then for each x in \mathcal{I}

$$f(x) = f(a) + f'(a)(x-a) + \frac{f''(a)}{2}(x-a)^2 + \frac{f'''(a)}{3!}(x-a)^3 + \cdots + \frac{f^{(n)}(a)}{n!}(x-a)^n + \frac{f^{(n+1)}(z)}{(n+1)!}(x-a)^{n+1}$$

where z is some number between a and x .

Let's revisit our first example and improve the approximation:

Example 4.1

Let $f(x) = x + \ln x$.

- (i) Find $T_3(x)$, the Taylor polynomial of degree 3 for f at $a = 1$.
- (ii) Use $T_3(x)$ to approximate $f(1.1)$.
- (iii) Give an error bound for your approximation.

Solution

- (i) This time

$$f(x) = x + \ln x ,$$

$$f'(x) = 1 + \frac{1}{x} ,$$

$$f''(x) = \frac{-1}{x^2}$$

$$\text{and } f'''(x) = \frac{2}{x^3}.$$

Evaluating these at $a = 1$ gives

$$f(1) = 1, \quad f'(1) = 2, \quad f''(1) = -1 \quad \text{and} \quad f'''(1) = 2$$

so we have

$$\begin{aligned} T_3(x) &= f(a) + f'(a)(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \frac{f'''(a)}{3!}(x-a)^3 \\ &= 1 + 2(x-1) - \frac{1}{2!}(x-1)^2 + \frac{2}{3!}(x-1)^3 \\ &= 1 + 2(x-1) - \frac{1}{2}(x-1)^2 + \frac{1}{3}(x-1)^3 \end{aligned}$$

(ii) The approximation is

$$\begin{aligned} f(1.1) &\approx T_3(1.1) \\ &= 1 + 2(1.1 - 1) - \frac{1}{2}(1.1 - 1)^2 + \frac{1}{3}(1.1 - 1)^3 \\ &= 1 + \frac{1}{5} - \frac{1}{200} + \frac{1}{3000} \\ &= \frac{1793}{1500} \end{aligned}$$

(iii) We require a bound on $|R_3(1.1)|$. By Theorem 4.1

$$R_3(1.1) = \frac{f^{(4)}(z)}{4!}(1.1 - 1)^4$$

for some $1 < z < 1.1$. Here $f^{(4)}(z) = -6/z^4$, so

$$|R_4(1.1)| = \left| \frac{-6}{4!z^4}(1.1 - 1)^4 \right| = \left| \frac{-1}{4z^4}(0.1)^4 \right|.$$

Now ask: what is the largest possible size of $|R_3(1.1)|$ for $1 < z < 1.1$? Again in this case, taking z as small as possible makes $|R_3(1.1)|$ as large as possible, so setting $z = 1$ we have

$$|R_3(1.1)| \leq \left| \frac{-1}{4(1)^4}(0.1)^4 \right| = \frac{1}{40,000} = 0.000025.$$

That is,

$$|f(1.1) - T_3(1.1)| \leq 0.000025.$$

The same comments about error bound determination listed in Example 3.1 apply here:

1. It is absolute value of the error that is of interest.
2. In the case of an increasing or decreasing error function, the error bound will be found as z approaches one of the endpoints of the interval.
3. For a general error bound function the bound need not occur at the endpoints of the interval; more sophisticated optimization techniques may be required.

□

Section 4 Exercises

- 4.1** Determine the Taylor polynomial of degree 3 about $a = -1$ for $f(x) = \cos(2 + 2x)$ and use it to approximate $f(-3/2)$. Give an error bound for your approximation.
- 4.2** (a) Find $T_1(x)$ and $T_2(x)$ for $f(x) = x^2 \ln(x)$ at $a = 1$.
- (b) If $T_2(x)$ is used to approximate $f(4/5)$ determine an error bound on the resulting approximation. (Note that you are not being asked to find the approximation to $f(4/5)$, but only the error bound associated with the approximation.)

- 4.3 Find $T_1(x)$ and $T_2(x)$ for $f(x) = (1+x)^k$ at $a = 0$ (k is a constant here).
- 4.4 Use $T_2(x)$ from the previous exercise to approximate $\sqrt[3]{1.009}$ and give an error bound for your approximation. (Hint: What must be x and k so that $(1+x)^k = \sqrt[3]{1.009}$?)
- 4.5 Find the Taylor polynomial of degree 3 for $f(x) = \tan x$ at $a = \pi/4$.
- 4.6 Use a Maclaurin polynomial of degree 3 to approximate $e^{-0.1}$ and give an error bound on your approximation.
- 4.7 Find the Maclaurin polynomial of degree 3 for $f(x) = xe^{-x}$.
- 4.8 Find the Maclaurin polynomials of degree 1, 2, 3 and 4 for $f(x) = 2 + 3x - 5x^2 - 7x^3 + 11x^4$.
- What do you notice?
- 4.9 Find the Taylor polynomial of degree 3 for $f(x) = 3 + x + 4x^2 - 2x^3$ at $a = 2$, and then expand and simplify your result. What do you notice?
- 4.10 (a) Find the Maclaurin polynomial of degree 5 for $f(x) = \sin x$.
- (b) Determine the value of b such that your approximation in part (a) is accurate to within 0.00005 for x in the interval $(-b, b)$. [This one is tricky: here $T_5(x) = T_6(x)$ (why?) so b can be determined by analyzing either $R_5(x)$ or $R_6(x)$, each resulting in a different b value.]
- 4.11 Find the Maclaurin polynomial of degree 4 for $f(x) = e^x$. Based on your work, what would be the Maclaurin polynomial of degree n ?

Answers

- 4.1 $T_3(x) = 1 - 2(x+1)^2$; $f(-3/2) \approx T_3(-3/2) = 1/2$;
 $|R_3(-3/2)| \leq 1/24$
- 4.2 (a) $T_1(x) = x - 1$; $T_2(x) = (x-1) + \frac{3(x-1)^2}{2}$
 (b) $|R_2(4/5)| \leq 1/300$
- 4.3 $T_1(x) = 1 + kx$; $T_2(x) = 1 + kx + \frac{k(k-1)}{2}x^2$
- 4.4 $T_2(0.009) = \frac{1,002,991}{1,000,000}$; $|R_2(0.009)| \leq \frac{9}{2 \cdot 10^8} = 4.5 \times 10^{-8}$
- 4.5 $T_3(x) = 1 + 2\left(x - \frac{\pi}{4}\right) + 2\left(x - \frac{\pi}{4}\right)^2 + \frac{8}{3}\left(x - \frac{\pi}{4}\right)^3$
- 4.6 $T_3(x) = 1 + x + \frac{x^2}{2} + \frac{x^3}{6}$; $e^{-0.1} \approx T_3(-1/10) = 5429/6000$;
 $|R_3(-1/10)| \leq 1/240,000$
- 4.7 $T_3(x) = x - x^2 + \frac{x^3}{2}$
- 4.8 $T_1(x) = 2 + 3x$; $T_2(x) = 2 + 3x - 5x^2$; $T_3(x) = 2 + 3x - 5x^2 - 7x^3$;
 $T_4(x) = 2 + 3x - 5x^2 - 7x^3 + 11x^4$.
 Notice: for a polynomial $f(x)$, the Maclaurin polynomial of degree k simply consists of all terms of $f(x)$ up to and including that of degree k .
- 4.9 $T_3(x) = 5 - 7(x-2) - 8(x-2)^2 - 2(x-2)^3$.
 Notice: once expanded, $T_3(x) = f(x)$.
- 4.10 (a) $T_5(x) = x - \frac{x^3}{3!} + \frac{x^5}{5!}$
- (b) If analyzing $R_5(x)$, $b = (9/250)^{\frac{1}{6}}$. If analyzing $R_6(x)$,
 $b = (63/250)^{\frac{1}{7}}$.
- 4.11 $T_4(x) = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!}$;
 $T_n(x) = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \dots + \frac{x^n}{n!}$

5 Taylor & Maclaurin Series

In the previous section we saw that if f has $n+1$ derivatives on an open interval \mathcal{I} containing a then

$$f(x) = T_n(x) + R_n(x)$$

for every x in \mathcal{I} . This brings up a natural question: as n increases to ∞ , can $f(x)$ be somehow expressed as a polynomial of "infinite degree". The answer is, yes, but with conditions.

Definition 5.1 If f has derivatives of all orders at a then the **infinite series** (or simply **series**)

$$\begin{aligned} T(x) &= f(a) + f'(a)(x-a) + \frac{f''(a)}{2}(x-a)^2 + \frac{f'''(a)}{3!}(x-a)^3 + \frac{f^{(4)}(a)}{4!}(x-a)^4 + \cdots \\ &= \sum_{k=0}^{\infty} \frac{f^{(k)}(a)}{k!}(x-a)^k \end{aligned}$$

is called **the Taylor series for f at a** . In the case where $a = 0$ this series is called the **Maclaurin series for f** .¹

So if a function has derivatives of all orders at a then we can define its Taylor series $T(x)$, but is $f(x) = T(x)$? The answer is given by the following theorem:

Theorem 5.1 Suppose f has derivatives of all orders on an open interval \mathcal{I} containing a and that for each x in \mathcal{I}

$$\lim_{n \rightarrow \infty} R_n(x) = 0.$$

Then for each x in \mathcal{I} we have

$$f(x) = \sum_{k=0}^{\infty} \frac{f^{(k)}(a)}{k!}(x-a)^k$$

and we say that the Taylor (or Maclaurin) series $T(x)$ **converges** to $f(x)$ on \mathcal{I} . The largest open interval \mathcal{I} containing a on which $T(x)$ converges to $f(x)$ is called the **open interval of convergence** of the series.

To be precise, when $f(x)$ is equal to its Taylor series on an interval \mathcal{I} and we write

$$f(x) = \sum_{k=0}^{\infty} \frac{f^{(k)}(a)}{k!}(x-a)^k$$

we mean $f(x) = \lim_{n \rightarrow \infty} T_n(x)$ for each x in \mathcal{I}

A measure of caution is necessary when working with Taylor (or Maclaurin) series: although a function f may have derivatives of all orders at a , so that the Taylor series $T(x)$ is defined, it does not necessarily follow that $f(x) = T(x)$ when $x \neq a$. Furthermore, it is possible that a function is equal to its Taylor series on an interval but not on the entire domain of f . The analysis of $R_n(x)$ as $n \rightarrow \infty$ is critical in the determination of when a function is equal to its Taylor (or Maclaurin) series.

It turns out that many familiar functions are equal to their Taylor (or Maclaurin) series. Before looking at these we first state a limit result which often arises in the analysis of $R_n(x)$ (the proof is given in the Appendix):

Theorem 5.2 For each real number x ,

$$\lim_{n \rightarrow \infty} \frac{|x|^n}{n!} = 0$$

¹Note the introduction of **sigma notation** \sum here which is convenient for representing the sum of a large number of terms, particularly when the terms follow a pattern. See the appendix in the back of *Essential Calculus, Early Transcendentals* by James Stewart (1st or 2nd edition) for a review of sigma notation.

As a first example, let's find the Maclaurin series for $f(x) = e^x$ and show that it converges to e^x for every real number x . Because this is such an important Maclaurin series we state the result as a theorem:

Theorem 5.3 For every real number x ,

$$e^x = \sum_{k=0}^{\infty} \frac{x^k}{k!} = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \cdots$$

Proof

Every derivative of e^x is again e^x , so with $a = 0$ we have $f(0) = f'(0) = f''(0) = \cdots = f^{(n)}(0) = 1$. The resulting Maclaurin series for e^x is

$$\begin{aligned} T(x) &= f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3 + \frac{f^{(4)}(0)}{4!}x^4 + \cdots \\ &= 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \cdots \\ &= \sum_{k=0}^{\infty} \frac{x^k}{k!}. \end{aligned}$$

To show that this Maclaurin series is equal to e^x for every real number x , consider the error associated with the Maclaurin polynomial of degree n :

$$R_n(x) = \frac{f^{(n+1)}(z)}{(n+1)!}x^{n+1} = \frac{e^z}{(n+1)!}x^{n+1} \text{ where } z \text{ is some number between } 0 \text{ and } x.$$

We must show that for each x , $R_n(x) \rightarrow 0$. If $x = 0$ then $R_n(x) = 0$ and we're done. If $x \neq 0$,

$$\begin{aligned} |R_n(x)| &= \left| \frac{e^z x^{n+1}}{(n+1)!} \right| \\ &= \frac{e^z |x|^{n+1}}{(n+1)!} \text{ since } e^z > 0 \\ &\leq \frac{e^{|z|} |x|^{n+1}}{(n+1)!} \text{ since } z \leq |z| \\ &\leq \frac{e^{|x|} |x|^{n+1}}{(n+1)!} \text{ since } |z| < |x| \end{aligned}$$

$\rightarrow 0$ as $n \rightarrow \infty$ by Theorem 5.2.

Since $|R_n(x)| \rightarrow 0$ so does $R_n(x)$, and so by Theorem 5.1

$$e^x = \sum_{k=0}^{\infty} \frac{x^k}{k!}$$

□

Showing that $R_n(x) \rightarrow 0$ is not always so easy and there are other methods for showing that a function is equal to its Taylor or Maclaurin series. Here is a second example, this time for a function which is not equal to its Maclaurin series on its entire domain, and we'll use a different method to analyze $R_n(x)$:

Example 5.1

Find the Maclaurin series for $f(x) = \frac{1}{1-x}$ and determine the open interval of convergence.

Solution

First note that the domain of f is all real numbers excluding $x = 1$. Evaluating f and its derivatives at $a = 0$ we find

$$f(x) = \frac{1}{(1-x)} \quad ; \quad f(0) = 1$$

$$f'(x) = \frac{1}{(1-x)^2} \quad ; \quad f'(0) = 1$$

$$f''(x) = \frac{2}{(1-x)^3} \quad ; \quad f''(0) = 2$$

$$f'''(x) = \frac{3!}{(1-x)^4} \quad ; \quad f'''(0) = 3!$$

$$f^{(4)}(x) = \frac{4!}{(1-x)^5} \quad ; \quad f^{(4)}(0) = 4!$$

and in general,

$$f^{(k)}(x) = \frac{k!}{(1-x)^{k+1}} \quad ; \quad f^{(k)}(0) = k! .$$

Inserting this into the formula for the Maclaurin series we find

$$\begin{aligned} T(x) &= 1 + 1 \cdot x + \frac{2}{2}x^2 + \frac{3!}{3!}x^3 + \frac{4!}{4!}x^4 + \dots \\ &= 1 + x + x^2 + x^3 + x^4 + \dots \\ &= \sum_{k=0}^{\infty} x^k . \end{aligned}$$

To analyze $R_n(x)$ we'll use a method that is unique to this example. For any x in the domain of f , (i.e. $x \neq 1$),

$$T_n(x) = 1 + x + x^2 + x^3 + x^4 + \dots + x^n .$$

Multiplying both sides of this expression by x gives

$$xT_n(x) = x + x^2 + x^3 + x^4 + \cdots + x^{n+1} .$$

Subtracting the second equation from the first then gives

$$\begin{aligned} T_n(x) - xT_n(x) &= 1 + x + x^2 + x^3 + x^4 + \cdots + x^n \\ &\quad - x - x^2 - x^3 - x^4 - \cdots - x^n - x^{n+1} . \end{aligned}$$

Factoring on the left and cancelling on the right results in

$$(1 - x)T_n(x) = 1 - x^{n+1} ,$$

and isolating $T_n(x)$ then gives

$$T_n(x) = \frac{1 - x^{n+1}}{1 - x} .$$

The error term can then be expressed

$$R_n(x) = f(x) - T_n(x) = \frac{1}{1 - x} - \frac{1 - x^{n+1}}{1 - x} = \frac{x^{n+1}}{1 - x} .$$

Notice here that $R_n(x)$ is the exact error on the domain of f and does not depend on some unknown z between 0 and x . Furthermore, we see that $R_n(x) \rightarrow 0$ as $n \rightarrow \infty$ if and only if $|x| < 1$, i.e. $-1 < x < 1$. Therefore,

$$\frac{1}{1 - x} = \sum_{k=0}^{\infty} x^k \text{ for } x \text{ in } (-1, 1) .$$

□

The series

$$\frac{1}{1 - x} = \sum_{k=0}^{\infty} x^k \text{ for } x \text{ in } (-1, 1)$$

from the previous example is called the **geometric series** and it is useful for constructing new Taylor series from existing ones as we'll see in the next section. Other standard series are likewise useful in this respect so we list them in a table of standard Maclaurin series at the end of this section.

Another important observation about Taylor (and Maclaurin) series is that they contain the Taylor (*resp.* Maclaurin) polynomials of every degree for the function in question. For example, using the fact that

$$\ln(1 + x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \cdots \text{ on } (-1, 1)$$

we can immediately conclude that, for example, the Maclaurin polynomial of degree 5 for $f(x) = \ln(1 + x)$ is

$$T_5(x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \frac{x^5}{5} .$$

Maclaurin Series	Open Interval of Convergence
$e^x = \sum_{k=0}^{\infty} \frac{x^k}{k!} = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots$	$(-\infty, \infty)$
$\sin x = \sum_{k=0}^{\infty} \frac{(-1)^k x^{2k+1}}{(2k+1)!} = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots$	$(-\infty, \infty)$
$\cos x = \sum_{k=0}^{\infty} \frac{(-1)^k x^{2k}}{(2k)!} = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots$	$(-\infty, \infty)$
$\ln(1+x) = \sum_{k=0}^{\infty} \frac{(-1)^k x^{k+1}}{k+1} = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots$	$(-1, 1)$
$\frac{1}{1-x} = \sum_{k=0}^{\infty} x^k = 1 + x + x^2 + x^3 + \dots$	$(-1, 1)$
$\arctan x = \sum_{k=0}^{\infty} \frac{(-1)^k x^{2k+1}}{2k+1} = x - \frac{x^3}{3} + \frac{x^5}{5} - \dots$	$(-1, 1)$

Section 5 Exercises

5.1 Show that $\sin x$ is equal to its Maclaurin series on $(-\infty, \infty)$.

5.2 Find the Maclaurin series for $f(x) = 2 + 3x - 5x^2 - 7x^3 + 11x^4$ and state the open interval of convergence. This example shows that a Maclaurin (or Taylor) series need not have an infinite number of non-zero terms.

5.3 Find the Maclaurin series for $f(x) = xe^x$.

5.4 Find the Taylor series for $f(x) = 3 + x + 4x^2 - 2x^3$ at $a = 2$ and state the open interval of convergence.

5.5 Find the Maclaurin series for $f(x) = \frac{x^2}{x+1}$ (Hint: to make the calculation of derivatives easier, use

long division of polynomials to first express $f(x)$ in the form

$$f(x) = ax + b + \frac{c}{x+1}$$

for suitable a , b and c .)

5.6 Find the Taylor series at $a = 1$ for $f(x) = \frac{1}{x^2}$.

5.7 What is the sum of the infinite series

$$1 + 1 + \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!} + \dots ?$$

5.8 What is the sum of the infinite series

$$\pi - \frac{\pi^3}{3!} + \frac{\pi^5}{5!} - \dots ?$$

5.9 The Maclaurin series for $f(x) = x^3 e^{-x^2}$ is

$$T(x) = \sum_{k=0}^{\infty} (-1)^k \frac{x^{2k+3}}{k!}.$$

Determine $f^{(13)}(0)$ and $f^{(14)}(0)$.

5.10 Use the fact that

$$\frac{1}{1-x} = \sum_{k=0}^{\infty} x^k \text{ on } (-1, 1)$$

to find a Maclaurin series which is equal to $g(x) = \frac{1}{1+x^2}$. What is the open interval of convergence?

Answers

5.2 $T(x) = 2 + 3x - 5x^2 - 7x^3 + 11x^4; (-\infty, \infty)$

5.3 $T(x) = \sum_{k=0}^{\infty} \frac{x^{k+1}}{k!}$

5.4 $T(x) = 5 - 7(x-2) - 8(x-2)^2 - 2(x-2)^3; (-\infty, \infty)$

5.5 $T(x) = \sum_{k=0}^{\infty} (-1)^k x^{k+2}$

5.6 $T(x) = \sum_{k=0}^{\infty} (-1)^k (k+1)(x-1)^k$

5.7 $e^1 = e$

5.8 $\sin \pi = 0$

5.9 $f^{(13)}(0) = -\frac{13!}{5!} = 51,891,840$ and $f^{(14)}(0) = 0$

5.10 $T(x) = \sum_{k=0}^{\infty} (-1)^k x^{2k}; \mathcal{I} = (-1, 1)$

6 Working with Taylor & Maclaurin Series

Functions that are equal to their Taylor (*resp.* Maclaurin) series are equal to the limit as $n \rightarrow \infty$ of their Taylor (*resp.* Maclaurin) polynomials. This fact, together with standard limit laws, allows us to construct Taylor and Maclaurin series for new functions by combining the Taylor and Maclaurin series of known functions in much the same way one would treat polynomials.

Theorem 6.1 *Suppose*

$$f(x) = \sum_{k=0}^{\infty} b_k (x-a)^k$$

and

$$g(x) = \sum_{k=0}^{\infty} c_k (x-a)^k$$

are Taylor series representations for f and g at the point a with open intervals of convergence \mathcal{I}_1 and \mathcal{I}_2 , respectively. Then on $\mathcal{I} = \mathcal{I}_1 \cap \mathcal{I}_2$, (that is, \mathcal{I} is the largest open interval contained in both \mathcal{I}_1 and \mathcal{I}_2),

(i) $(f+g)(x) = \sum_{k=0}^{\infty} p_k (x-a)^k$ where $p_k = b_k + c_k$

(ii) $(f-g)(x) = \sum_{k=0}^{\infty} p_k (x-a)^k$ where $p_k = b_k - c_k$

$$(iii) (fg)(x) = \sum_{k=0}^{\infty} p_k(x-a)^k \text{ where } p_k = \sum_{j=0}^k b_j c_{k-j}$$

For each of the new series in (i), (ii) and (iii), the open interval of convergence may be \mathcal{I} itself or possibly a larger interval containing \mathcal{I} .

In simpler terms, this theorem tells us that the Taylor series for the sum, difference and product of two functions is found by simply adding, subtracting, and multiplying (respectively) the Taylor series for the two functions as one would do with polynomials. That this is true for sums and differences of functions is not that surprising. The fact that this is also true for products is less obvious and is given in the Appendix. The reason that Theorem 6.1 is true has to do with a property of more general series we'll look at in the next section. For now we quote the essential fact:

Theorem 6.2 *Suppose the series*

$$f(x) = r_0 + r_1(x-a) + r_2(x-a)^2 + r_3(x-a)^3 + \cdots = \sum_{k=0}^{\infty} r_k(x-a)^k$$

converges for every x in an open interval \mathcal{I} containing a . Then f is differentiable on \mathcal{I} ,

$$f'(x) = r_1 + 2r_2(x-a) + 3r_3(x-a)^2 + 4r_4(x-a)^3 + \cdots = \sum_{k=1}^{\infty} k r_k(x-a)^{k-1}$$

and this series for $f'(x)$ also converges on \mathcal{I} .

This theorem tells us that the Taylor (or Maclaurin) series for the derivative of a function is obtained by simply differentiating the Taylor (*resp.* Maclaurin) series of the original function term by term.

Here are some examples illustrating the application of these theorems:

Example 6.1

Find the Maclaurin series for $f(x) = x^2 e^x$.

Solution

$$\begin{aligned} x^2 e^x &= x^2 \left(1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots \right) \\ &= x^2 + x^3 + \frac{x^4}{2!} + \frac{x^5}{3!} + \cdots \\ &= \sum_{k=0}^{\infty} \frac{x^{k+2}}{k!} \end{aligned}$$

Since the Maclaurin series for x^2 and e^x both converge on $(-\infty, \infty)$, the open interval of convergence for x^2e^x will again be $(-\infty, \infty)$. \square

Notice in this example that the final answer was stated using \sum notation. This is not strictly necessary, but is good practice in the case where the terms of the sum follow an obvious pattern. You will generally be told the form to use for expressing your final answer.

Example 6.2

Find the first four non-zero terms of the Maclaurin series for $f(x) = e^x \cos x$.

Solution

Multiply each term from the first series by each term in the second and collect like terms:

$$\begin{aligned} e^x \cos x &= \left(1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots\right) \left(1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots\right) \\ &= 1 \cdot \left(1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots\right) + x \cdot \left(1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots\right) \\ &\quad + \frac{x^2}{2!} \cdot \left(1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots\right) + \frac{x^3}{3!} \cdot \left(1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots\right) \\ &\quad + \frac{x^4}{4!} \cdot \left(1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots\right) + \dots \\ &= 1 + x - \frac{x^3}{3} - \frac{x^4}{6} + \dots \end{aligned}$$

Again in this example, the Maclaurin series for e^x and $\cos x$ both converge on $(-\infty, \infty)$ and so the open interval of convergence for the product is also $(-\infty, \infty)$. \square

Another way to define new functions from old is by function composition. Recall that $(f \circ g)(x) = f(g(x))$. In the case where $g(x) = x^r$ for r a non-negative integer we have

Theorem 6.3

$$f(x) = \sum_{k=0}^{\infty} b_k x^k$$

is a Maclaurin series for f with open interval of convergence \mathcal{I} and r is a non-negative integer such that x^r is in \mathcal{I} , then

$$f(x^r) = \sum_{k=0}^{\infty} b_k x^{rk}$$

is the Maclaurin series for $f(x^r)$.

Example 6.3

Find the Maclaurin polynomial of degree 6 for $f(x) = \frac{x}{1 + 4x^2}$.

Solution

Let's find the Maclaurin series for f and then extract the terms up to degree 6. First rearrange $f(x)$ to make the choice of series more obvious:

$$f(x) = \frac{x}{1 + 4x^2} = x \cdot \frac{1}{1 - (-4x^2)}.$$

For the fraction $\frac{1}{1 - (-4x^2)}$ we'll use the geometric series for $\frac{1}{1 - x}$ with x replaced with $-4x^2$:

$$\begin{aligned} f(x) &= x \cdot \frac{1}{1 - (-4x^2)} \\ &= x \cdot (1 + (-4x^2) + (-4x^2)^2 + (-4x^2)^3 + (-4x^2)^4 + \dots) \\ &= x \cdot (1 - 4x^2 + 16x^4 - 64x^6 + \dots) \\ &= x - 4x^3 + 16x^5 - 64x^7 + \dots \end{aligned}$$

Extracting the terms of degree 6 or less we have

$$T_6(x) = x - 4x^3 + 16x^5.$$

□

Taylor and Maclaurin series can be used to calculate limits, and in some cases this approach is far more efficient than L'Hospital's Rule as the following example illustrates:

Example 6.4

Evaluate $\lim_{x \rightarrow 0} \frac{e^{-x^2} + x^2 - 1}{x^4}$

Solution

Applying direct substitution yields the indeterminate form " $\frac{0}{0}$ ". Let's replace e^{-x^2} with its

Maclaurin series, which is easily found by replacing x with $-x^2$ in the series for e^x :

$$\begin{aligned} \lim_{x \rightarrow 0} \frac{e^{-x^2} + x^2 - 1}{x^4} &= \lim_{x \rightarrow 0} \frac{\left(1 + (-x^2) + \frac{(-x^2)^2}{2!} + \frac{(-x^2)^3}{3!} + \dots\right) + x^2 - 1}{x^4} \\ &= \lim_{x \rightarrow 0} \frac{\left(1 - x^2 + \frac{x^4}{2!} - \frac{x^6}{3!} + \dots\right) + x^2 - 1}{x^4} \\ &= \lim_{x \rightarrow 0} \frac{\left(\frac{x^4}{2!} - \frac{x^6}{3!} + \dots\right)}{x^4} \\ &= \lim_{x \rightarrow 0} \frac{x^4 \left(\frac{1}{2!} - \frac{x^2}{3!} + \dots\right)}{x^4} \\ &= \lim_{x \rightarrow 0} \left(\frac{1}{2!} - \frac{x^2}{3!} + \dots\right) \\ &= \frac{1}{2} \end{aligned}$$

□

Section 6 Exercises

- 6.1** Find the Maclaurin series and state the open interval of convergence:
- $f(x) = \cos(\pi x)$
 - $f(x) = x \arctan x$
- 6.2** Find the first four non-zero terms of the Maclaurin series for $f(x) = e^x \ln(1 - x)$.
- 6.3** Find the first four non-zero terms of the Maclaurin series for $f(x) = \frac{\sin x}{1 + x}$.
- 6.4** Use a series to evaluate the following limits:
- $\lim_{x \rightarrow 0} \frac{1 - \cos x}{1 + x - e^x}$
 - $\lim_{x \rightarrow 0} \frac{\sin x - x + \frac{x^3}{6}}{x^5}$
 - $\lim_{x \rightarrow 0} \frac{\ln(1 + x^2) - e^{(x^2)} + 1}{2x^4}$
- 6.5** Find the Maclaurin polynomial of degree 13 for $f(x) = x^3 \sin(x^2)$.
- 6.6** Find the Maclaurin series for $f(x) = \frac{x}{(1+x)^2}$
(Hint: $\frac{d}{dx} \left[\frac{1}{1+x} \right] = \frac{-1}{(1+x)^2}$.)
- 6.7** Find the Maclaurin series and state the open interval of convergence for $f(x) = \sin^2 x$. (Hint: using a trigonometric identity to first rewrite $f(x)$ will make this easier.)
- 6.8** Find the Maclaurin series for $g(x) = 2 \sin x \cos x$ (Hint: use your result from the previous exercise.)
- 6.9** The hyperbolic sine function, $\sinh x$, is defined to be
- $$\sinh x = \frac{e^x - e^{-x}}{2}.$$
- This function arises in the solution to certain equations in physics and engineering. Find the Maclaurin series for $\sinh(x^2/3)$.

6.10 We know that $\frac{d}{dx}[e^x] = e^x$. Prove this by differentiating the Maclaurin series for e^x .

6.11 Show that the derivative of $\sin x$ is $\cos x$ using Maclaurin series.

6.12 It can be shown that the Maclaurin series for $\sin^4 x$ is

$$\sin^4 x = x^4 - \frac{2x^6}{3} + \frac{x^8}{5} - \dots$$

Use this to find the first three nonzero terms of the Maclaurin series for $g(x) = 4 \sin^3(x) \cos(x)$.

6.13 Find the Taylor series for $f(x) = \frac{1}{7-3x}$

(a) about $a = 1$

(b) about $a = -1$

and state the open interval of convergence for each. (This one is tricky: think about using the series for $1/(1-x)$.)

6.14 Find the the first four nonzero terms of the Taylor series for $h(x) = \frac{1}{2-3x}$ about $a = 1$ and state the open interval of convergence.

6.15 Use a Maclaurin series to prove that $\lim_{x \rightarrow 0} \frac{\sin(x)}{x} = 1$.

Answers

6.1 (a) $\sum_{k=0}^{\infty} \frac{(-1)^k \pi^{2k} x^{2k}}{(2k)!}; (-\infty, \infty)$

(b) $\sum_{k=0}^{\infty} \frac{(-1)^k x^{2(k+1)}}{2k+1}; (-1, 1)$

6.2 $-x - \frac{3x^2}{2} - \frac{4x^3}{3} - x^4 - \dots$

6.3 $x - x^2 + \frac{5x^3}{6} - \frac{5x^4}{6} + \dots$

6.4 (a) -1

(b) $\frac{1}{120}$

(c) $-\frac{1}{2}$

6.5 $T_{13}(x) = x^5 - \frac{x^9}{6} + \frac{x^{13}}{120}$

6.6 $\sum_{k=0}^{\infty} (-1)^k (k+1)x^{k+1}$

6.7 $\sum_{k=0}^{\infty} \frac{(-1)^k 2^{2k+1} x^{2(k+1)}}{(2(k+1))!}; (-\infty, \infty)$

6.8 $\sum_{k=0}^{\infty} \frac{(-1)^k 2^{2k+1} x^{2k+1}}{(2k+1)!}$

6.9 $\sum_{k=0}^{\infty} \frac{x^{2(2k+1)}}{3^{2k+1}(2k+1)!}$

6.12 $4x^3 - 4x^5 + \frac{8x^7}{5} + \dots$

6.13 (a) $\sum_{k=0}^{\infty} \left(\frac{1}{4}\right) \left(\frac{3}{4}\right)^k (x-1)^k; \mathcal{I} = (-1/3, 7/3)$

(b) $\sum_{k=0}^{\infty} \left(\frac{1}{10}\right) \left(\frac{3}{10}\right)^k (x+1)^k; \mathcal{I} = (-13/3, 7/3)$

6.14 $-1 + 3(x-1) - 9(x-1)^2 + 27(x-1)^3 + \dots; \mathcal{I} = (2/3, 4/3)$

7 Power Series

So far we have seen that if a function f has derivatives of all orders at a point $x = a$ then the Taylor (or Maclaurin) series for the function can be defined, and under suitable conditions, the series converges to the function on an open interval of convergence \mathcal{I} . That is, for x in \mathcal{I} ,

$$f(x) = \sum_{k=0}^{\infty} a_k (x-a)^k,$$

where equality here is understood to mean

$$f(x) = \lim_{n \rightarrow \infty} \sum_{k=0}^n a_k (x - a)^k .$$

Now let's examine this idea from the other direction: suppose we have an infinite sequence of real numbers a_0, a_1, a_2, \dots and we form the infinite series

$$\sum_{k=0}^{\infty} a_k (x - a)^k .$$

Now let

$$f(x) = \sum_{k=0}^{\infty} a_k (x - a)^k = \lim_{n \rightarrow \infty} \sum_{k=0}^n a_k (x - a)^k ,$$

where the domain of f is the set of real numbers x for which the limit above exists. What can we say about functions defined in this way? In particular, what is the domain, is the function differentiable, and if so, what is the derivative?

Definition 7.1 A series of the form

$$\sum_{k=0}^{\infty} a_k (x - a)^k = a_0 + a_1(x - a) + a_2(x - a)^2 + a_3(x - a)^3 + \dots$$

is called a **power series centred at a** or a **power series about a** . The real numbers a_0, a_1, a_2, \dots are called the **coefficients** of the series.

Notice that Taylor and Maclaurin series fall into the general category of power series. Also notice that $f(a) = a_0$ for every power series $f(x) = \sum_{k=0}^{\infty} a_k (x - a)^k$, so there is at least one point in the domain of f . To determine the open interval of convergence of the power series, we turn to the following theorem which is a consequence of a result called the **ratio test** from the general theory of infinite series:

Theorem 7.1 Suppose $\sum_{k=0}^{\infty} a_k (x - a)^k$ is a power series and let $u_k(x)$ be the k^{th} non-zero term of the series (note that some of the a_k may be zero). Then the power series converges if

$$\lim_{k \rightarrow \infty} \frac{|u_{k+1}(x)|}{|u_k(x)|} < 1$$

We illustrate the use of the theorem with two examples.

Example 7.1

Determine the open interval of convergence of

$$\sum_{k=0}^{\infty} \frac{(-1)^k x^{k+1}}{k+1}$$

Solution

Here $u_k(x) = \frac{(-1)^k x^{k+1}}{k+1}$. For convergence we require

$$\lim_{k \rightarrow \infty} \frac{|u_{k+1}(x)|}{|u_k(x)|} < 1$$

$$\lim_{k \rightarrow \infty} \frac{\left| \frac{(-1)^{k+1} x^{k+2}}{k+2} \right|}{\left| \frac{(-1)^k x^{k+1}}{k+1} \right|} < 1$$

$$\lim_{k \rightarrow \infty} \left| \frac{x^{k+2}}{x^{k+1}} \frac{k+1}{k+2} \right| < 1$$

$$\lim_{k \rightarrow \infty} \left| x \frac{k+1}{k+2} \right| < 1$$

$$|x| < 1$$

So the open interval of convergence is $-1 < x < 1$, i.e. $\mathcal{I} = (-1, 1)$.

You may notice from our work on Taylor series that the power series in this example is the Taylor series defining $\ln(1+x)$, and $\mathcal{I} = (-1, 1)$ is the result stated in the summary of Maclaurin series at the end of Section 5. \square

Example 7.2

Determine the open interval of convergence of $\sum_{k=1}^{\infty} \frac{(-1)^k 4^k (x-1)^{2k}}{k}$.

Solution

Here

$$u_k(x) = \frac{(-1)^k 4^k (x-1)^{2k}}{k}$$

For convergence of the series we require

$$\lim_{k \rightarrow \infty} \frac{|u_{k+1}(x)|}{|u_k(x)|} < 1$$

$$\lim_{k \rightarrow \infty} \left| \frac{\frac{(-1)^{k+1} 4^{k+1} (x-1)^{2(k+1)}}{k+1}}{\frac{(-1)^k 4^k (x-1)^{2k}}{k}} \right| < 1$$

$$\lim_{k \rightarrow \infty} \left| \frac{4^{k+1} (x-1)^{2k+2}}{4^k (x-1)^{2k}} \frac{k}{k+1} \right| < 1$$

$$\lim_{k \rightarrow \infty} \left| 4(x-1)^2 \frac{k}{k+1} \right| < 1$$

$$4|x-1|^2 < 1$$

$$|x-1|^2 < \frac{1}{4}$$

$$\text{so } |x-1| < \frac{1}{2}$$

That is, the series converges for $1/2 < x < 3/2$, so the open interval of convergence is $\mathcal{I} = (1/2, 3/2)$. \square

In these two examples, the limit

$$\lim_{k \rightarrow \infty} \frac{|u_{k+1}(x)|}{|u_k(x)|} < 1$$

resulted in a statement of the form $|x-a| < R$ which gave the open interval of convergence $(a-R, a+R)$. The value R here is called the **radius of convergence** of the power series. There are three possibilities for R :

- (i) If $R = 0$, the power series converges only for $x = a$ and the open interval of convergence does not exist.
- (ii) If $R > 0$, the power series converges for $|x-a| < R$ and diverges for $|x-a| > R$. That is, the open interval of convergence of the power series is $(a-R, a+R)$.
- (iii) If $R = \infty$ the power series converges for all real numbers: the open interval of convergence is $(-\infty, \infty)$.

The first two examples above show possibility (ii) where $R > 0$ is a finite number. The next two examples show the other two possibilities: $R = 0$ and $R = \infty$:

Example 7.3

[$(R = 0)$] Determine the radius of convergence and open interval of convergence of

$$\sum_{k=0}^{\infty} k!(x-2)^k$$

Solution

Here $u_k(x) = k!(x-2)^k$. We require

$$\begin{aligned} \lim_{k \rightarrow \infty} \frac{|u_{k+1}(x)|}{|u_k(x)|} &< 1 \\ \lim_{k \rightarrow \infty} \frac{|(k+1)!(x-2)^{k+1}|}{|k!(x-2)^k|} &< 1 \\ \lim_{k \rightarrow \infty} |(k+1)(x-2)| &< 1 \end{aligned}$$

which is only possible if $x = 2$, and so $R = 0$. In this case the open interval of convergence does not exist. \square

Example 7.4

[$(R = \infty)$] Determine the radius of convergence and open interval of convergence of

$$\sum_{k=0}^{\infty} \frac{(-1)^k x^{2k}}{2^{2k} (k!)^2}$$

Solution

Here $u_k = \frac{(-1)^k x^{2k}}{2^{2k} (k!)^2}$ and we require

$$\lim_{k \rightarrow \infty} \frac{|u_{k+1}(x)|}{|u_k(x)|} < 1$$

$$\lim_{k \rightarrow \infty} \frac{\left| \frac{(-1)^{k+2} x^{2(k+1)}}{2^{2(k+1)} ((k+1)!)^2} \right|}{\left| \frac{(-1)^k x^{2k}}{2^{2k} (k!)^2} \right|} < 1$$

$$\lim_{k \rightarrow \infty} \left| \frac{2^{2k}}{2^{2k+2}} \frac{k!}{(k+1)!} \frac{k!}{(k+1)!} \frac{x^{2k+2}}{x^{2k}} \right| < 1$$

$$\lim_{k \rightarrow \infty} \left| \frac{1}{4} \frac{1}{(k+1)^2} x^2 \right| < 1$$

which results in $0 < 1$,

and this last statement is true for every real number x . So $R = \infty$ in this case and the open interval of convergence is the entire real number line, $\mathcal{I} = (-\infty, \infty)$.

(The function defined by the power series in this example is called a Bessel function of order 0. Bessel functions arise in the solutions of problems involving wave propagation, among others.) \square

In the case where $R > 0$ and finite, the power series may also converge for $x = a - R$ or for $x = a + R$ (that is, at one or both of the boundary points of the open interval of convergence.) The convergence behaviour at each of these two points must be analyzed separately using other techniques from the general theory of infinite series. For our purposes, it will suffice to find the radius of convergence and, in the case of $R > 0$, the corresponding open interval of convergence.

Derivatives of Power Series

The derivatives of functions defined by power series can be found as one might expect, by differentiating term by term as one would a polynomial. The details are contained the in the following theorem which we state without proof:

Theorem 7.2 Suppose $\sum_{k=0}^{\infty} a_k(x-a)^k$ is a power series with radius of convergence $R > 0$ and corresponding open interval of convergence $\mathcal{I} = (a - R, a + R)$. Then the function

$$f(x) = \sum_{k=0}^{\infty} a_k(x-a)^k$$

is differentiable on \mathcal{I} ,

$$\begin{aligned} f'(x) &= a_1 + 2a_2(x-a) + 3a_3(x-a)^2 + 4a_4(x-a)^3 + \cdots \\ &= \sum_{k=1}^{\infty} k a_k (x-a)^{k-1}, \end{aligned}$$

and the radius of convergence of the power series defining $f'(x)$ is again R .

As a consequence of this theorem, if $f(x) = \sum_{k=0}^{\infty} a_k (x-a)^k$ on \mathcal{I} , then

$$a_k = \frac{f^{(k)}(a)}{k!} \text{ for } k = 0, 1, 2, \dots$$

In other words, the power series $\sum_{k=0}^{\infty} a_k (x-a)^k$ is the Taylor series about a for the function f it defines, and the open interval of convergence of the Taylor series is \mathcal{I} .

Example 7.5

The first five non-zero terms of the power series representing $\tan x$ on $\mathcal{I} = (-\pi/2, \pi/2)$ are

$$\tan x = x + \frac{x^3}{3} + \frac{2x^5}{15} + \frac{17x^7}{315} + \frac{62x^9}{2835} + \cdots$$

Use this information to find the Maclaurin series for $\sec^2 x$. State the open interval of convergence.

Solution

Since the given series represents $\tan x$ on \mathcal{I} , it is equal to the Maclaurin series for the function. Differentiating gives

$$\begin{aligned} \frac{d}{dx} [\tan x] &= \frac{d}{dx} \left[x + \frac{x^3}{3} + \frac{2x^5}{15} + \frac{17x^7}{315} + \frac{62x^9}{2835} + \cdots \right] \\ \sec^2 x &= 1 + \frac{3x^2}{3} + \frac{10x^4}{15} + \frac{119x^6}{315} + \frac{558x^8}{2835} + \cdots \\ &= 1 + x^2 + \frac{2x^4}{3} + \frac{17x^6}{45} + \frac{62x^8}{315} + \cdots \end{aligned}$$

The open interval of convergence is the same as that of the original series: $\mathcal{I} = (-\pi/2, \pi/2)$ \square

Section 7 Exercises

7.1 Find the radius of convergence and open interval of convergence:

$$(a) \sum_{k=0}^{\infty} \frac{(x-2)^k}{10^k}$$

$$(b) \sum_{k=0}^{\infty} \frac{(3x-2)^{k+1}}{k+1} \quad [\text{caution: put the series in powers of } (x-a) \text{ first}]$$

$$(c) \sum_{k=1}^{\infty} \frac{(x-1)^k}{k^3 3^k}$$

$$(d) \sum_{k=1}^{\infty} \frac{(-1)^k 3^{2k} (x-2)^k}{3k}$$

$$(e) \sum_{k=0}^{\infty} k!(x-4)^k$$

$$(f) \sum_{k=0}^{\infty} \frac{(-1)^k (x-1)^{4k}}{k!}$$

$$(g) \sum_{k=0}^{\infty} \frac{(-1)^k (x-3)^{2k}}{9^k}$$

$$(h) \sum_{k=1}^{\infty} \frac{4^k (x-3)^{2k}}{k^2}$$

$$(i) \sum_{k=0}^{\infty} \frac{(-1)^k (x+1)^k}{2^k k!}$$

$$(j) \sum_{k=1}^{\infty} \frac{k^k x^k}{k!} \quad (\text{Hint: } \lim_{k \rightarrow \infty} (1 + \frac{1}{k})^k = e)$$

7.2 Find the radius of convergence and open interval of convergence of

$$\sum_{k=0}^{\infty} \frac{x^k}{e^k}.$$

What function does this series represent?

7.3 Use the result in Example 7.5 to find $f^{(8)}(0)$ for $f(x) = \tan^2 x$. (Hint: use the identity $1 + \tan^2 x = \sec^2 x$.)

Answers

7.1 (a) $R = 10; \mathcal{I} = (-8, 12)$

(b) $R = 1/3; \mathcal{I} = (1/3, 1)$

(c) $R = 3; \mathcal{I} = (-2, 4)$

(d) $R = 1/9; \mathcal{I} = (17/9, 19/9)$

(e) $R = 0; \mathcal{I}$ does not exist (series converges only for $x = 4$)

(f) $R = \infty; \mathcal{I} = (-\infty, \infty)$

(g) $R = 3; \mathcal{I} = (0, 6)$

(h) $R = 1/2; \mathcal{I} = (5/2, 7/2)$

(i) $R = \infty; \mathcal{I} = (-\infty, \infty)$

(j) $R = 1/e; \mathcal{I} = (-1/e, 1/e)$

7.2 $R = e; \mathcal{I} = (-e, e); f(x) = e/(e-x)$

7.3 7936

Appendix

Theorem 3.1 Suppose that f is twice differentiable on an open interval \mathcal{I} containing containing a . Then for each x in \mathcal{I}

$$R_1(x) = \frac{f''(z)}{2}(x - a)^2$$

for some z between a and x .

Proof

The proof uses Rolle's theorem applied to a specially constructed function. Suppose a and x are in \mathcal{I} , $x \neq a$, and consider x fixed (that is, a constant). Define

$$g(t) = f(x) - f(t) - f'(t)(x - t) - R_1(x) \frac{(x - t)^2}{(x - a)^2}.$$

Notice that $g(a) = g(x) = 0$, so by Rolle's Theorem there is some number z strictly between a and x such that $g'(z) = 0$. Now

$$\begin{aligned} g'(t) &= 0 - f'(t) - f''(t)(x - t) + f'(t) + 2R_1(x) \frac{(x - t)}{(x - a)^2} \\ &= -f''(t)(x - t) + 2R_1(x) \frac{(x - t)}{(x - a)^2} \end{aligned}$$

so

$$g'(z) = -f''(z)(x - z) + 2R_1(x) \frac{(x - z)}{(x - a)^2} = 0$$

giving

$$R_1(x) = \frac{f''(z)}{2}(x - a)^2$$

□

Theorem 4.1 Suppose f is $(n + 1)$ -times differentiable on an open interval \mathcal{I} containing containing a . Then for each x in \mathcal{I}

$$R_n(x) = \frac{f^{(n+1)}(z)}{(n + 1)!}(x - a)^{n+1}$$

for some z between a and x .

Proof

The proof is a generalization of the $n = 1$ case above. Suppose a and x are in \mathcal{I} , $x \neq a$, and consider x fixed. Define

$$g(t) = f(x) - f(t) - f'(t)(x - t) - \frac{f''(t)}{2!}(x - t)^2 - \dots - \frac{f^{(n)}(t)}{n!}(x - t)^n - R_n(x) \frac{(x - t)^{n+1}}{(x - a)^{n+1}}$$

and notice that $g(a) = g(x) = 0$. By Rolle's Theorem there is some number z strictly between a and x such that $g'(z) = 0$. Now differentiating and simplifying we find

$$g'(t) = -\frac{f^{(n+1)}(t)}{n!}(x-t)^n + (n+1)R_n(x)\frac{(x-t)^n}{(x-a)^{(n+1)}}$$

so that

$$g'(z) = -\frac{f^{(n+1)}(z)}{n!}(x-z)^n + (n+1)R_n(x)\frac{(x-z)^n}{(x-a)^{(n+1)}}.$$

Setting $g'(z) = 0$ and isolating $R_n(x)$ then gives

$$R_n(x) = \frac{f^{(n+1)}(z)}{(n+1)!}(x-a)^{n+1}.$$

□

Theorem 5.2 For each real number x ,

$$\lim_{n \rightarrow \infty} \frac{|x|^n}{n!} = 0$$

Proof

Let x be any fixed real number. If $0 \leq |x| < 1$ then

$$\frac{|x|^n}{n!} \leq \frac{1}{n!} \rightarrow 0 \text{ as } n \rightarrow \infty.$$

If $|x| \geq 1$, let m be the unique positive integer such that $m-1 \leq |x| < m$. Then

$$\begin{aligned} \frac{|x|^n}{n!} &= \frac{|x|}{1} \cdot \frac{|x|}{2} \cdots \frac{|x|}{n} \\ &= \frac{|x|}{1} \cdots \frac{|x|}{m-1} \left(\frac{|x|}{m} \cdot \frac{|x|}{m+1} \cdots \frac{|x|}{n} \right) \\ &\leq \frac{|x|^{m-1}}{(m-1)!} \left(\frac{|x|}{m} \right)^{n-m+1} \\ &\rightarrow 0 \text{ as } n \rightarrow \infty \text{ since } \frac{|x|}{m} < 1 \end{aligned}$$

In either case, $\lim_{n \rightarrow \infty} \frac{|x|^n}{n!} = 0$.

□

Theorem 6.1 *Suppose*

$$f(x) = \sum_{k=0}^{\infty} b_k(x-a)^k$$

and

$$g(x) = \sum_{k=0}^{\infty} c_k(x-a)^k$$

are Taylor series representations for f and g at the point a with open intervals of convergence \mathcal{I}_1 and \mathcal{I}_2 , respectively. Then on $\mathcal{I} = \mathcal{I}_1 \cap \mathcal{I}_2$, (that is, \mathcal{I} is the largest open interval contained in both \mathcal{I}_1 and \mathcal{I}_2),

$$(i) (f+g)(x) = \sum_{k=0}^{\infty} p_k(x-a)^k \text{ where } p_k = b_k + c_k$$

$$(ii) (f-g)(x) = \sum_{k=0}^{\infty} p_k(x-a)^k \text{ where } p_k = b_k - c_k$$

$$(iii) (fg)(x) = \sum_{k=0}^{\infty} p_k(x-a)^k \text{ where } p_k = \sum_{j=0}^k b_j c_{k-j}$$

For each of the new series in (i), (ii) and (iii), the open interval of convergence may be \mathcal{I} itself or possibly a larger interval containing \mathcal{I} .

Proof

The proofs of (i), (ii) and (iii) are similar; let's prove (iii) since it's the least obvious of the three. Let $F_n(x)$ and $G_n(x)$ be the Taylor polynomials of degree n at a for f and g , respectively. By hypothesis

$$f(x) = \lim_{n \rightarrow \infty} F_n(x) \text{ for every } x \text{ in } \mathcal{I}_1$$

and

$$g(x) = \lim_{n \rightarrow \infty} G_n(x) \text{ for every } x \text{ in } \mathcal{I}_2 .$$

Since these limits exist and \mathcal{I} is contained in both \mathcal{I}_1 and \mathcal{I}_2 , then for every x in \mathcal{I}

$$\begin{aligned} f(x)g(x) &= \left(\lim_{n \rightarrow \infty} F_n(x) \right) \left(\lim_{n \rightarrow \infty} G_n(x) \right) \\ &= \lim_{n \rightarrow \infty} (F_n(x)G_n(x)) \\ &= \sum_{k=0}^{\infty} p_k(x-a)^k \end{aligned}$$

where $p_k = \sum_{j=0}^k b_j c_{k-j}$ are the coefficients resulting from multiplying the series together and collecting like terms.

So $(fg)(x)$ has a convergent series representation on \mathcal{I} ; to complete the proof we must show that this series is indeed a Taylor series. That is, we must show that for each k ,

$$p_k = \frac{1}{k!} \frac{d^k}{dx^k} [f(x)g(x)]_{x=a} .$$

Setting $x = a$ in

$$f(x)g(x) = p_0 + p_1(x - a) + p_2(x - a)^2 + p_3(x - a)^3 + p_4(x - a)^4 + \cdots$$

we find

$$p_0 = f(a)g(a) .$$

Taking k derivatives and setting $x = a$ in

$$f(x)g(x) = p_0 + p_1(x - a) + p_2(x - a)^2 + p_3(x - a)^3 + p_4(x - a)^4 + \cdots$$

gives (by Theorem 6.2)

$$\begin{aligned} \frac{d^k}{dx^k} [f(x)g(x)]_{x=a} &= \frac{d^k}{dx^k} [p_0 + p_1(x - a) + p_2(x - a)^2 + p_3(x - a)^3 + p_4(x - a)^4 + \cdots]_{x=a} \\ &= [k!p_k + (\text{terms with at least one factor of } (x - a))]_{x=a} \\ &= k!p_k \end{aligned}$$

so that

$$p_k = \frac{1}{k!} \frac{d^k}{dx^k} [f(x)g(x)]_{x=a} ,$$

which completes the proof. □