

## 1 General Derivative Rules

1. Constant Rule  $\frac{d}{dx} [c] = 0$
2. Constant Multiple Rule  $\frac{d}{dx} [cf(x)] = cf'(x)$
3. Sum Rule  $\frac{d}{dx} [f(x) + g(x)] = f'(x) + g'(x)$
4. Difference Rule  $\frac{d}{dx} [f(x) - g(x)] = f'(x) - g'(x)$
5. Product Rule  $\frac{d}{dx} [f(x)g(x)] = f'(x)g(x) + f(x)g'(x)$
6. Quotient Rule  $\frac{d}{dx} \left[ \frac{f(x)}{g(x)} \right] = \frac{g(x)f'(x) - f(x)g'(x)}{[g(x)]^2}$
7. Chain Rule  $\frac{d}{dx} [f(g(x))] = f'(g(x))g'(x)$

## 2 Derivative Rules for Particular Functions

	Basic Rule	Chain Rule Form
1. Powers	$\frac{d}{dx} [x^n] = nx^{n-1}$	$\frac{d}{dx} [(f(x))^n] = n(f(x))^{n-1}f'(x)$
2. Sine	$\frac{d}{dx} [\sin x] = \cos x$	$\frac{d}{dx} [\sin (f(x))] = \cos (f(x))f'(x)$
3. Cosine	$\frac{d}{dx} [\cos x] = -\sin x$	$\frac{d}{dx} [\cos (f(x))] = -\sin (f(x))f'(x)$
4. Tangent	$\frac{d}{dx} [\tan x] = \sec^2 x$	$\frac{d}{dx} [\tan (f(x))] = \sec^2 (f(x))f'(x)$
5. Secant	$\frac{d}{dx} [\sec x] = \sec x \tan x$	$\frac{d}{dx} [\sec (f(x))] = \sec (f(x)) \tan (f(x))f'(x)$
6. Cosecant	$\frac{d}{dx} [\csc x] = -\csc x \cot x$	$\frac{d}{dx} [\csc (f(x))] = -\csc (f(x)) \cot (f(x))f'(x)$
7. Cotangent	$\frac{d}{dx} [\cot x] = -\csc^2 x$	$\frac{d}{dx} [\cot (f(x))] = -\csc^2 (f(x))f'(x)$
8. Exponential (base e)	$\frac{d}{dx} [e^x] = e^x$	$\frac{d}{dx} [e^{(f(x))}] = e^{(f(x))}f'(x)$
9. Exponential (base a)	$\frac{d}{dx} [a^x] = a^x \ln a$	$\frac{d}{dx} [a^{(f(x))}] = a^{(f(x))} \ln a f'(x)$
10. Natural Logarithm	$\frac{d}{dx} [\ln x] = \frac{1}{x}$	$\frac{d}{dx} [\ln f(x)] = \frac{1}{f(x)}f'(x)$
11. Logarithm (base a)	$\frac{d}{dx} [\log_a x] = \frac{1}{x \ln a}$	$\frac{d}{dx} [\log_a f(x)] = \frac{1}{f(x) \ln a} f'(x)$
12. Inverse sine	$\frac{d}{dx} [\arcsin x] = \frac{1}{\sqrt{1-x^2}}$	$\frac{d}{dx} [\arcsin f(x)] = \frac{1}{\sqrt{1-(f(x))^2}} f'(x)$
13. Inverse cosine	$\frac{d}{dx} [\arccos x] = \frac{-1}{\sqrt{1-x^2}}$	$\frac{d}{dx} [\arccos f(x)] = \frac{-1}{\sqrt{1-(f(x))^2}} f'(x)$
14. Inverse tangent	$\frac{d}{dx} [\arctan x] = \frac{1}{1+x^2}$	$\frac{d}{dx} [\arctan f(x)] = \frac{1}{1+(f(x))^2} f'(x)$

### 3 General Antiderivative Rules

Let  $F(x)$  be any antiderivative of  $f(x)$ . That is,  $F'(x) = f(x)$ . The most general antiderivative of  $f(x)$  is then  $F(x) + C$ .

	Original Function	General Antiderivative
1. Constant Rule	$c$ (a constant)	$cx + C$
2. Constant Multiple Rule	$cf(x)$	$cF(x) + C$
3. Sum Rule	$f(x) + g(x)$	$F(x) + G(x) + C$
4. Difference Rule	$f(x) - g(x)$	$F(x) - G(x) + C$

### 4 Antiderivative Rules for Particular Functions

	Original Function	General Antiderivative
1. Powers ( $n \neq -1$ )	$x^n$	$\frac{x^{n+1}}{n+1} + C$
2. Powers ( $n = -1$ )	$\frac{1}{x}$	$\ln x  + C$
3. Sine	$\sin x$	$-\cos x + C$
4. Cosine	$\cos x$	$\sin x + C$
5. Secant squared	$\sec^2 x$	$\tan x + C$
6. Secant times tangent	$\sec x \tan x$	$\sec x + C$
7. Cosecant times cotangent	$\csc x \cot x$	$-\csc x + C$
8. Cosecant squared	$\csc^2 x$	$-\cot x + C$
9. Exponential (base $e$ )	$e^x$	$e^x + C$
10. Exponential (base $a$ )	$a^x$	$\frac{a^x}{\ln a} + C$
11. Inverse trigonometric	$\frac{1}{\sqrt{1-x^2}}$	$\arcsin x + C$
12. Inverse trigonometric	$\frac{-1}{\sqrt{1-x^2}}$	$\arccos x + C$
13. Inverse trigonometric	$\frac{1}{1+x^2}$	$\arctan x + C$